

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex**Session Type:** Rate**Suggested Citation****Software:** Surveillance Research Program, National Cancer Institute SEER*Stat software (www.seer.cancer.gov/seerstat) version 6.5.2.**Data:** Alaska Cancer Registry SEER*Stat Database - Incidence data for diagnosis years 1996 to 2006. Released May 2009. Alaska population data for rate calculations are from the SEER program (<http://www.seer.cancer.gov/popdata/>) and include populations by borough/census area, expanded races, and Hispanic origin.**Data****Database:** ACR w/Borough, Race, & Hispanic Pops (1996-2006), May 2009**Statistic****Statistic:** Age-Adjusted Rates (Age recode with <1 year olds) (2000 US Std Population (19 age groups - Census P25-1130) standard)**P-Values:** .05**Options:** Display Confidence Intervals (Tiwari mod) and Standard Errors**Selection****Select Only:** Malignant Behavior, Known Age, Male or Female Sex**Table****Page:** Year of diagnosis**Row:** Site rec with Kaposi and mesothelioma**Column:** Race recode (4 Races - W, B, AI, API) [Race recode (W, B, AI, API)]
Sex - M & F only [Sex]**Output****Title:** Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex**Suppress:** Statistics based on fewer than 6 cases**Display Rates as:** Cases per 100,000**Precision:** 1 decimal place (default)

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 561.5 | 548.6 | 574.7 | 9,653 | 434.5 | 424.4 | 444.7 | 8,180 |
| Oral Cavity and Pharynx | 15.2 | 13.3 | 17.2 | 318 | 6.3 | 5.1 | 7.6 | 107 |
| Lip | 1.7 | 1.0 | 2.7 | 26 | 0.6 | 0.2 | 1.1 | 9 |
| Tongue | 4.0 | 3.1 | 5.1 | 87 | 1.4 | 0.9 | 2.0 | 25 |
| Salivary Gland | 1.5 | 0.9 | 2.3 | 23 | 0.9 | 0.5 | 1.5 | 15 |
| Floor of Mouth | 1.0 | 0.6 | 1.5 | 22 | 0.5 | 0.2 | 1.0 | 9 |
| Gum and Other Mouth | 1.6 | 1.0 | 2.4 | 32 | 1.0 | 0.6 | 1.7 | 17 |
| Nasopharynx | 0.7 | 0.4 | 1.2 | 18 | 0.4 | 0.2 | 0.9 | 8 |
| Tonsil | 2.7 | 2.0 | 3.5 | 69 | 0.4 | 0.2 | 0.8 | 7 |
| Oropharynx | 0.7 | 0.4 | 1.3 | 15 | 0.4 | 0.2 | 0.9 | 7 |
| Hypopharynx | 0.9 | 0.5 | 1.5 | 18 | 0.5 | 0.2 | 1.0 | 8 |
| Other Oral Cavity and Pharynx | 0.3 | 0.1 | 0.7 | 8 | ^ | ^ | ^ | ^ |
| Digestive System | 100.8 | 95.2 | 106.6 | 1,670 | 69.3 | 65.0 | 73.7 | 1,120 |
| Esophagus | 8.9 | 7.4 | 10.6 | 163 | 2.2 | 1.6 | 3.1 | 38 |
| Stomach | 7.8 | 6.2 | 9.5 | 121 | 3.0 | 2.2 | 4.1 | 50 |
| Small Intestine | 1.9 | 1.3 | 2.7 | 36 | 1.5 | 0.9 | 2.3 | 25 |
| Colon and Rectum | 56.7 | 52.5 | 61.1 | 911 | 40.6 | 37.3 | 44.0 | 645 |
| Colon excluding Rectum | 41.4 | 37.7 | 45.2 | 618 | 29.9 | 27.1 | 32.9 | 458 |
| Cecum | 9.0 | 7.2 | 11.0 | 122 | 9.1 | 7.6 | 10.8 | 133 |
| Appendix | 0.6 | 0.3 | 1.1 | 17 | ^ | ^ | ^ | ^ |
| Ascending Colon | 7.6 | 6.0 | 9.4 | 103 | 5.6 | 4.4 | 7.1 | 78 |
| Hepatic Flexure | 1.7 | 1.1 | 2.5 | 29 | 1.8 | 1.2 | 2.6 | 27 |
| Transverse Colon | 2.6 | 1.8 | 3.5 | 44 | 1.9 | 1.2 | 2.7 | 28 |
| Splenic Flexure | 1.0 | 0.5 | 1.7 | 14 | 0.9 | 0.4 | 1.5 | 13 |
| Descending Colon | 3.6 | 2.6 | 4.8 | 52 | 1.1 | 0.7 | 1.8 | 21 |
| Sigmoid Colon | 11.7 | 9.8 | 13.8 | 182 | 6.9 | 5.6 | 8.4 | 115 |
| Large Intestine, NOS | 3.7 | 2.6 | 5.0 | 55 | 2.3 | 1.6 | 3.2 | 38 |
| Rectum and Rectosigmoid Junction | 15.3 | 13.4 | 17.5 | 293 | 10.6 | 9.1 | 12.4 | 187 |
| Rectosigmoid Junction | 4.1 | 3.1 | 5.3 | 73 | 2.1 | 1.4 | 2.9 | 37 |
| Rectum | 11.2 | 9.6 | 13.0 | 220 | 8.6 | 7.2 | 10.2 | 150 |
| Anus, Anal Canal and Anorectum | 1.0 | 0.5 | 1.7 | 19 | 2.8 | 2.1 | 3.8 | 56 |
| Liver and Intrahepatic Bile Duct | 8.5 | 7.0 | 10.2 | 158 | 3.2 | 2.4 | 4.2 | 58 |
| Liver | 7.8 | 6.3 | 9.5 | 145 | 2.6 | 1.8 | 3.5 | 46 |
| Intrahepatic Bile Duct | 0.7 | 0.3 | 1.3 | 13 | 0.6 | 0.3 | 1.1 | 12 |
| Gallbladder | 0.9 | 0.4 | 1.6 | 13 | 0.9 | 0.5 | 1.6 | 12 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 647.2 | 569.9 | 730.9 | 363 | 371.1 | 319.9 | 427.4 | 249 |
| Oral Cavity and Pharynx | 9.3 | 2.5 | 23.3 | 6 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Tonsil | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Digestive System | 155.8 | 116.1 | 203.0 | 81 | 74.0 | 50.8 | 103.0 | 40 |
| Esophagus | 12.0 | 3.7 | 27.7 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 24.1 | 10.4 | 45.6 | 10 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 68.0 | 42.0 | 101.9 | 38 | 58.4 | 37.8 | 84.8 | 30 |
| Colon excluding Rectum | 46.5 | 26.4 | 74.1 | 28 | 46.2 | 28.2 | 70.1 | 24 |
| Cecum | 6.2 | 2.2 | 16.1 | 6 | 9.8 | 3.2 | 22.0 | 6 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | 13.6 | 4.7 | 28.9 | 6 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 8.2 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 12.9 | 4.3 | 28.0 | 6 |
| Large Intestine, NOS | 15.9 | 3.8 | 38.4 | 7 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 21.5 | 7.5 | 44.8 | 10 | 12.2 | 4.0 | 26.5 | 6 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Rectum | 12.0 | 3.7 | 27.7 | 6 | 12.2 | 4.0 | 26.5 | 6 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 27.0 | 11.6 | 50.9 | 13 | ^ | ^ | ^ | ^ |
| Liver | 27.0 | 11.6 | 50.9 | 13 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

Total

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 535.0 | 507.0 | 564.1 | 1,655 | 500.1 | 477.0 | 524.0 | 1,907 |
| Oral Cavity and Pharynx | 22.8 | 17.9 | 28.5 | 82 | 11.3 | 8.0 | 15.3 | 43 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Tongue | 3.2 | 1.6 | 5.8 | 12 | 2.3 | 1.0 | 4.5 | 9 |
| Salivary Gland | 1.5 | 0.5 | 3.3 | 6 | 1.6 | 0.7 | 3.3 | 8 |
| Floor of Mouth | 2.3 | 1.0 | 4.5 | 8 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 1.7 | 0.6 | 3.8 | 6 | ^ | ^ | ^ | ^ |
| Nasopharynx | 9.3 | 6.2 | 13.3 | 32 | 4.5 | 2.5 | 7.4 | 16 |
| Tonsil | 1.3 | 0.5 | 3.1 | 6 | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Digestive System | 199.7 | 182.2 | 218.2 | 586 | 158.2 | 144.8 | 172.4 | 554 |
| Esophagus | 12.8 | 8.8 | 17.8 | 38 | 8.1 | 5.2 | 11.8 | 26 |
| Stomach | 30.5 | 24.1 | 37.9 | 96 | 15.8 | 11.9 | 20.5 | 61 |
| Small Intestine | 2.2 | 0.6 | 4.9 | 6 | 2.1 | 0.9 | 4.1 | 8 |
| Colon and Rectum | 107.6 | 94.9 | 121.3 | 314 | 99.8 | 89.3 | 111.2 | 348 |
| Colon excluding Rectum | 77.1 | 66.2 | 89.2 | 214 | 77.4 | 68.0 | 87.5 | 265 |
| Cecum | 12.7 | 8.5 | 18.1 | 34 | 18.6 | 14.2 | 23.9 | 62 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Ascending Colon | 8.6 | 5.2 | 13.1 | 23 | 9.6 | 6.4 | 13.6 | 31 |
| Hepatic Flexure | 4.9 | 2.4 | 8.6 | 14 | 5.3 | 3.0 | 8.5 | 17 |
| Transverse Colon | 9.4 | 5.7 | 14.3 | 24 | 12.7 | 9.0 | 17.4 | 39 |
| Splenic Flexure | 3.7 | 1.6 | 7.0 | 9 | 3.0 | 1.5 | 5.3 | 12 |
| Descending Colon | ^ | ^ | ^ | ^ | 3.1 | 1.4 | 5.6 | 10 |
| Sigmoid Colon | 32.2 | 25.5 | 39.9 | 95 | 22.5 | 17.8 | 27.9 | 85 |
| Large Intestine, NOS | 2.9 | 1.0 | 6.1 | 7 | 2.6 | 1.1 | 4.9 | 9 |
| Rectum and Rectosigmoid Junction | 30.4 | 24.3 | 37.5 | 100 | 22.5 | 17.7 | 28.1 | 83 |
| Rectosigmoid Junction | 7.7 | 4.9 | 11.5 | 25 | 4.6 | 2.6 | 7.3 | 17 |
| Rectum | 22.7 | 17.5 | 29.0 | 75 | 17.9 | 13.7 | 23.0 | 66 |
| Anus, Anal Canal and Anorectum | 2.4 | 1.1 | 4.6 | 10 | 2.3 | 1.0 | 4.4 | 8 |
| Liver and Intrahepatic Bile Duct | 12.6 | 8.4 | 17.9 | 38 | 6.0 | 3.7 | 9.0 | 23 |
| Liver | 11.3 | 7.3 | 16.4 | 34 | 4.9 | 2.9 | 7.8 | 19 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 5.6 | 2.8 | 9.7 | 13 | 4.8 | 2.6 | 7.9 | 15 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

Total

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 409.7 | 363.6 | 459.4 | 356 | 289.9 | 261.0 | 320.9 | 438 |
| Oral Cavity and Pharynx | 8.6 | 3.5 | 17.1 | 9 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Hypopharynx | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Digestive System | 125.9 | 102.1 | 153.1 | 118 | 65.2 | 51.1 | 81.8 | 87 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 25.0 | 14.3 | 39.5 | 19 | 8.5 | 4.1 | 15.3 | 11 |
| Small Intestine | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 43.5 | 30.0 | 60.5 | 39 | 27.2 | 18.2 | 38.6 | 35 |
| Colon excluding Rectum | 27.8 | 17.1 | 42.2 | 24 | 17.7 | 10.8 | 27.1 | 23 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 9.0 | 3.1 | 19.0 | 6 | 8.7 | 4.0 | 16.0 | 12 |
| Large Intestine, NOS | 6.7 | 2.5 | 14.1 | 7 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 15.7 | 8.4 | 26.5 | 15 | 9.5 | 4.5 | 17.2 | 12 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 9.8 | 4.5 | 18.5 | 10 | 6.5 | 2.4 | 13.5 | 8 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Liver and Intrahepatic Bile Duct | 32.0 | 21.9 | 45.2 | 37 | 13.8 | 8.1 | 21.8 | 19 |
| Liver | 31.5 | 21.4 | 44.6 | 36 | 13.3 | 7.7 | 21.2 | 18 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 1.6 | 1.0 | 2.6 | 24 | 1.3 | 0.8 | 2.1 | 21 |
| Pancreas | 11.6 | 9.8 | 13.7 | 193 | 10.9 | 9.2 | 12.7 | 171 |
| Retroperitoneum | 0.3 | 0.1 | 0.6 | 6 | 0.3 | 0.1 | 0.6 | 6 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.9 | 0.5 | 1.5 | 14 |
| Other Digestive Organs | 1.6 | 1.0 | 2.4 | 24 | 1.6 | 1.0 | 2.5 | 24 |
| Respiratory System | 93.5 | 88.2 | 99.0 | 1,498 | 65.9 | 61.8 | 70.2 | 1,071 |
| Nose, Nasal Cavity and Middle Ear | 0.9 | 0.5 | 1.5 | 16 | 0.3 | 0.1 | 0.7 | 7 |
| Larynx | 6.5 | 5.2 | 7.9 | 119 | 1.3 | 0.8 | 1.9 | 26 |
| Lung and Bronchus | 85.9 | 80.8 | 91.3 | 1,357 | 64.1 | 60.1 | 68.3 | 1,034 |
| Pleura | 0.0 | 0.0 | 0.2 | 0 | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.2 | 0.1 | 0.5 | 6 | ^ | ^ | ^ | ^ |
| Bones and Joints | 0.9 | 0.6 | 1.4 | 28 | 1.2 | 0.8 | 1.7 | 29 |
| Soft Tissue including Heart | 3.6 | 2.7 | 4.6 | 79 | 2.6 | 1.9 | 3.5 | 56 |
| Skin excluding Basal and Squamous | 17.6 | 15.6 | 19.9 | 378 | 13.1 | 11.6 | 14.8 | 293 |
| Melanoma of the Skin | 15.5 | 13.6 | 17.6 | 343 | 12.1 | 10.6 | 13.8 | 273 |
| Other Non-Epithelial Skin | 2.1 | 1.3 | 3.1 | 35 | 1.0 | 0.6 | 1.6 | 20 |
| Breast | 0.8 | 0.4 | 1.3 | 15 | 138.8 | 133.3 | 144.4 | 2,827 |
| Female Genital System | 0.0 | 0.0 | 0.2 | 0 | 48.1 | 45.0 | 51.4 | 991 |
| Cervix Uteri | 0.0 | 0.0 | 0.2 | 0 | 8.1 | 6.9 | 9.4 | 193 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 0.2 | 0 | 22.7 | 20.6 | 25.1 | 457 |
| Corpus Uteri | 0.0 | 0.0 | 0.2 | 0 | 21.9 | 19.8 | 24.2 | 442 |
| Uterus, NOS | 0.0 | 0.0 | 0.2 | 0 | 0.9 | 0.4 | 1.4 | 15 |
| Ovary | 0.0 | 0.0 | 0.2 | 0 | 12.8 | 11.2 | 14.6 | 260 |
| Vagina | 0.0 | 0.0 | 0.2 | 0 | 0.9 | 0.5 | 1.5 | 15 |
| Vulva | 0.0 | 0.0 | 0.2 | 0 | 2.6 | 1.9 | 3.5 | 48 |
| Other Female Genital Organs | 0.0 | 0.0 | 0.2 | 0 | 1.0 | 0.6 | 1.6 | 18 |
| Male Genital System | 175.4 | 168.3 | 182.7 | 2,986 | 0.0 | 0.0 | 0.2 | 0 |
| Prostate | 168.2 | 161.1 | 175.4 | 2,782 | 0.0 | 0.0 | 0.2 | 0 |
| Testis | 6.3 | 5.4 | 7.3 | 185 | 0.0 | 0.0 | 0.2 | 0 |
| Penis | 0.9 | 0.4 | 1.5 | 16 | 0.0 | 0.0 | 0.2 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.2 | 0 |
| Urinary System | 67.0 | 62.4 | 71.7 | 1,067 | 19.8 | 17.6 | 22.1 | 339 |
| Urinary Bladder | 47.7 | 43.7 | 51.9 | 695 | 9.8 | 8.2 | 11.5 | 157 |
| Kidney and Renal Pelvis | 18.2 | 16.1 | 20.6 | 358 | 9.3 | 7.9 | 10.9 | 171 |
| Ureter | 0.7 | 0.4 | 1.3 | 11 | 0.6 | 0.3 | 1.2 | 10 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Pancreas | 15.0 | 5.3 | 32.2 | 7 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Respiratory System | 108.3 | 77.7 | 145.7 | 55 | 50.9 | 32.9 | 74.4 | 29 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 8.2 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 97.6 | 69.0 | 132.9 | 51 | 47.9 | 30.3 | 71.0 | 27 |
| Pleura | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 112.2 | 86.5 | 142.5 | 86 |
| Female Genital System | 0.0 | 0.0 | 8.2 | 0 | 38.9 | 25.0 | 57.4 | 31 |
| Cervix Uteri | 0.0 | 0.0 | 8.2 | 0 | 8.6 | 3.8 | 17.5 | 9 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 8.2 | 0 | 18.6 | 8.5 | 34.1 | 12 |
| Corpus Uteri | 0.0 | 0.0 | 8.2 | 0 | 18.6 | 8.5 | 34.1 | 12 |
| Uterus, NOS | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Ovary | 0.0 | 0.0 | 8.2 | 0 | 8.4 | 3.4 | 17.9 | 8 |
| Vagina | 0.0 | 0.0 | 8.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Male Genital System | 266.7 | 219.9 | 319.8 | 149 | 0.0 | 0.0 | 5.8 | 0 |
| Prostate | 265.7 | 218.9 | 318.7 | 147 | 0.0 | 0.0 | 5.8 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Penis | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Urinary System | 36.8 | 19.9 | 61.1 | 18 | 26.1 | 12.2 | 46.6 | 11 |
| Urinary Bladder | 13.4 | 4.5 | 29.7 | 6 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 23.4 | 10.3 | 44.1 | 12 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

Total

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 4.5 | 2.2 | 8.1 | 12 | 3.9 | 2.0 | 6.6 | 13 |
| Pancreas | 19.0 | 13.7 | 25.4 | 51 | 14.0 | 10.2 | 18.6 | 48 |
| Retroperitoneum | 0.0 | 0.0 | 1.2 | 0 | 0.0 | 0.0 | 0.9 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 2.5 | 0.9 | 5.2 | 7 | ^ | ^ | ^ | ^ |
| Respiratory System | 122.7 | 109.5 | 136.9 | 364 | 80.1 | 70.6 | 90.5 | 270 |
| Nose, Nasal Cavity and Middle Ear | 2.0 | 0.6 | 4.5 | 6 | ^ | ^ | ^ | ^ |
| Larynx | 4.5 | 2.6 | 7.3 | 17 | 2.1 | 0.9 | 4.1 | 8 |
| Lung and Bronchus | 114.7 | 101.9 | 128.5 | 338 | 76.3 | 67.0 | 86.4 | 256 |
| Pleura | 0.0 | 0.0 | 1.2 | 0 | 0.0 | 0.0 | 0.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | 1.6 | 0.5 | 3.8 | 7 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 2.3 | 1.2 | 4.2 | 13 | 2.7 | 1.3 | 4.8 | 12 |
| Skin excluding Basal and Squamous | 2.8 | 1.2 | 5.4 | 9 | 2.4 | 1.1 | 4.5 | 10 |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | 1.8 | 0.8 | 3.7 | 8 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 1.2 | 0 | 135.9 | 124.6 | 148.0 | 561 |
| Female Genital System | 0.0 | 0.0 | 1.2 | 0 | 38.0 | 32.1 | 44.6 | 161 |
| Cervix Uteri | 0.0 | 0.0 | 1.2 | 0 | 9.8 | 7.1 | 13.2 | 47 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 1.2 | 0 | 15.3 | 11.6 | 19.7 | 61 |
| Corpus Uteri | 0.0 | 0.0 | 1.2 | 0 | 15.3 | 11.6 | 19.7 | 61 |
| Uterus, NOS | 0.0 | 0.0 | 1.2 | 0 | 0.0 | 0.0 | 0.9 | 0 |
| Ovary | 0.0 | 0.0 | 1.2 | 0 | 10.2 | 7.2 | 13.9 | 41 |
| Vagina | 0.0 | 0.0 | 1.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 1.2 | 0 | 1.5 | 0.5 | 3.3 | 6 |
| Other Female Genital Organs | 0.0 | 0.0 | 1.2 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 83.1 | 72.1 | 95.1 | 252 | 0.0 | 0.0 | 0.9 | 0 |
| Prostate | 74.8 | 64.3 | 86.4 | 215 | 0.0 | 0.0 | 0.9 | 0 |
| Testis | 5.1 | 3.4 | 7.5 | 30 | 0.0 | 0.0 | 0.9 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Urinary System | 50.0 | 41.7 | 59.4 | 152 | 19.6 | 15.1 | 24.9 | 70 |
| Urinary Bladder | 20.8 | 15.4 | 27.3 | 59 | 5.0 | 2.8 | 8.1 | 16 |
| Kidney and Renal Pelvis | 27.6 | 21.6 | 34.5 | 88 | 13.9 | 10.3 | 18.3 | 52 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

Total

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 13.9 | 6.5 | 25.0 | 11 | 6.9 | 2.6 | 14.0 | 8 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 63.3 | 45.6 | 85.0 | 50 | 45.7 | 33.8 | 60.2 | 56 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Lung and Bronchus | 59.2 | 42.1 | 80.3 | 47 | 43.7 | 32.0 | 57.9 | 53 |
| Pleura | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 3.8 | 0 | 67.0 | 54.6 | 81.4 | 114 |
| Female Genital System | 0.0 | 0.0 | 3.8 | 0 | 41.3 | 31.8 | 52.9 | 72 |
| Cervix Uteri | 0.0 | 0.0 | 3.8 | 0 | 8.5 | 4.9 | 14.0 | 17 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 3.8 | 0 | 18.9 | 12.4 | 27.5 | 32 |
| Corpus Uteri | 0.0 | 0.0 | 3.8 | 0 | 18.9 | 12.4 | 27.5 | 32 |
| Uterus, NOS | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Ovary | 0.0 | 0.0 | 3.8 | 0 | 10.7 | 6.0 | 17.6 | 17 |
| Vagina | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 123.4 | 97.4 | 153.3 | 92 | 0.0 | 0.0 | 2.7 | 0 |
| Prostate | 120.8 | 95.0 | 150.5 | 88 | 0.0 | 0.0 | 2.7 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Penis | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Urinary System | 25.7 | 14.9 | 40.4 | 20 | 12.2 | 6.4 | 20.6 | 16 |
| Urinary Bladder | 13.9 | 5.9 | 26.2 | 9 | 6.2 | 2.0 | 13.5 | 6 |
| Kidney and Renal Pelvis | 9.1 | 3.9 | 17.7 | 9 | 6.1 | 2.8 | 11.6 | 10 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Eye and Orbit | 0.9 | 0.6 | 1.4 | 23 | 0.5 | 0.2 | 0.9 | 9 |
| Brain and Other Nervous System | 7.9 | 6.6 | 9.4 | 177 | 6.3 | 5.2 | 7.5 | 129 |
| Brain | 7.7 | 6.4 | 9.1 | 170 | 6.0 | 4.9 | 7.2 | 123 |
| Cranial Nerves Other Nervous System | 0.3 | 0.1 | 0.6 | 7 | 0.3 | 0.1 | 0.6 | 6 |
| Endocrine System | 5.5 | 4.5 | 6.6 | 140 | 14.3 | 12.8 | 15.9 | 359 |
| Thyroid | 4.9 | 4.0 | 5.9 | 126 | 13.7 | 12.2 | 15.3 | 345 |
| Other Endocrine including Thymus | 0.6 | 0.3 | 1.1 | 14 | 0.6 | 0.3 | 1.1 | 14 |
| Lymphoma | 29.7 | 26.9 | 32.7 | 579 | 21.6 | 19.4 | 23.9 | 403 |
| Hodgkin Lymphoma | 3.9 | 3.1 | 4.9 | 99 | 2.2 | 1.6 | 2.9 | 53 |
| Hodgkin - Nodal | 3.9 | 3.1 | 4.9 | 99 | 2.1 | 1.6 | 2.8 | 52 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 0.2 | 0 | ^ | ^ | ^ | ^ |
| Non-Hodgkin Lymphoma | 25.7 | 23.1 | 28.6 | 480 | 19.4 | 17.3 | 21.7 | 350 |
| NHL - Nodal | 18.7 | 16.5 | 21.2 | 353 | 13.5 | 11.8 | 15.4 | 249 |
| NHL - Extranodal | 7.0 | 5.6 | 8.6 | 127 | 5.9 | 4.7 | 7.2 | 101 |
| Myeloma | 6.0 | 4.8 | 7.4 | 107 | 4.7 | 3.7 | 5.9 | 78 |
| Leukemia | 18.1 | 15.8 | 20.5 | 312 | 11.5 | 9.9 | 13.3 | 201 |
| Lymphocytic Leukemia | 10.5 | 8.8 | 12.4 | 183 | 5.8 | 4.6 | 7.1 | 99 |
| Acute Lymphocytic Leukemia | 1.8 | 1.2 | 2.5 | 41 | 1.1 | 0.7 | 1.6 | 28 |
| Chronic Lymphocytic Leukemia | 7.8 | 6.2 | 9.5 | 121 | 4.4 | 3.4 | 5.7 | 67 |
| Other Lymphocytic Leukemia | 1.0 | 0.5 | 1.6 | 21 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 6.6 | 5.2 | 8.1 | 112 | 4.8 | 3.8 | 6.0 | 86 |
| Acute Myeloid Leukemia | 3.9 | 2.9 | 5.2 | 66 | 2.7 | 1.9 | 3.6 | 48 |
| Acute Monocytic Leukemia | 0.2 | 0.1 | 0.6 | 6 | 0.4 | 0.1 | 0.8 | 7 |
| Chronic Myeloid Leukemia | 2.3 | 1.5 | 3.3 | 37 | 1.6 | 1.0 | 2.4 | 28 |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Leukemia | 1.0 | 0.5 | 1.7 | 17 | 0.9 | 0.5 | 1.5 | 16 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.7 | 0.3 | 1.4 | 12 | 0.7 | 0.3 | 1.3 | 12 |
| Mesothelioma | 4.7 | 3.4 | 6.3 | 55 | 0.5 | 0.2 | 0.9 | 7 |
| Kaposi Sarcoma | 0.2 | 0.1 | 0.5 | 7 | ^ | ^ | ^ | ^ |
| Miscellaneous | 13.8 | 11.7 | 16.1 | 214 | 10.2 | 8.6 | 12.0 | 159 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 8.2 | 0 | ^ | ^ | ^ | ^ |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 13.1 | 6.9 | 23.3 | 15 |
| Thyroid | ^ | ^ | ^ | ^ | 11.6 | 5.7 | 21.5 | 13 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 19.2 | 10.4 | 33.4 | 20 | 16.2 | 6.9 | 30.9 | 12 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Non-Hodgkin Lymphoma | 17.8 | 9.2 | 32.0 | 17 | 13.9 | 5.2 | 28.5 | 9 |
| NHL - Nodal | 9.5 | 3.8 | 21.0 | 10 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 8.3 | 2.8 | 20.0 | 7 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 9.9 | 2.6 | 24.6 | 6 | 9.8 | 3.8 | 20.5 | 9 |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 9.8 | 3.8 | 20.5 | 9 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 8.2 | 0 | ^ | ^ | ^ | ^ |
| Other Leukemia | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 8.2 | 0 | 0.0 | 0.0 | 5.8 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.8 | 0 |
| Miscellaneous | 14.2 | 5.3 | 30.0 | 9 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

Total

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Brain and Other Nervous System | 3.5 | 1.9 | 5.9 | 17 | 3.1 | 1.8 | 5.1 | 18 |
| Brain | 3.3 | 1.7 | 5.7 | 16 | 2.8 | 1.6 | 4.7 | 16 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 4.0 | 2.0 | 7.0 | 17 | 11.1 | 8.3 | 14.5 | 56 |
| Thyroid | 3.6 | 1.7 | 6.5 | 14 | 11.1 | 8.3 | 14.5 | 56 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Lymphoma | 15.3 | 11.1 | 20.4 | 56 | 12.7 | 9.3 | 16.8 | 53 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 1.2 | 0 | 0.0 | 0.0 | 0.9 | 0 |
| Non-Hodgkin Lymphoma | 14.4 | 10.3 | 19.5 | 52 | 12.3 | 9.0 | 16.4 | 51 |
| NHL - Nodal | 6.9 | 4.2 | 10.5 | 26 | 6.8 | 4.4 | 9.9 | 28 |
| NHL - Extranodal | 7.5 | 4.6 | 11.5 | 26 | 5.6 | 3.4 | 8.5 | 23 |
| Myeloma | 2.8 | 1.3 | 5.2 | 11 | 5.6 | 3.5 | 8.5 | 22 |
| Leukemia | 11.9 | 8.3 | 16.4 | 48 | 6.2 | 4.0 | 9.0 | 30 |
| Lymphocytic Leukemia | 2.7 | 1.3 | 5.0 | 15 | 2.7 | 1.4 | 4.7 | 14 |
| Acute Lymphocytic Leukemia | 1.3 | 0.7 | 2.8 | 11 | 1.6 | 0.7 | 3.1 | 10 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 1.2 | 0 | 0.0 | 0.0 | 0.9 | 0 |
| Myeloid and Monocytic Leukemia | 8.6 | 5.5 | 12.7 | 30 | 3.3 | 1.8 | 5.7 | 15 |
| Acute Myeloid Leukemia | 4.9 | 2.5 | 8.4 | 15 | 2.0 | 0.9 | 4.0 | 9 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | 2.8 | 1.3 | 5.3 | 11 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.9 | 0 |
| Miscellaneous | 10.5 | 6.8 | 15.2 | 31 | 11.8 | 8.3 | 16.1 | 41 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Asian or Pacific Islander | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Brain and Other Nervous System | 4.7 | 1.6 | 11.0 | 6 | 3.5 | 1.3 | 8.0 | 6 |
| Brain | 4.7 | 1.6 | 11.0 | 6 | 3.5 | 1.3 | 8.0 | 6 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 24.3 | 16.7 | 34.0 | 40 |
| Thyroid | ^ | ^ | ^ | ^ | 24.3 | 16.7 | 34.0 | 40 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Lymphoma | 10.9 | 5.4 | 19.6 | 13 | 11.4 | 6.6 | 18.5 | 18 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Non-Hodgkin Lymphoma | 10.9 | 5.4 | 19.6 | 13 | 10.5 | 5.8 | 17.4 | 16 |
| NHL - Nodal | 8.0 | 3.4 | 15.9 | 10 | 6.3 | 2.8 | 12.0 | 10 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 4.2 | 1.5 | 9.3 | 6 |
| Myeloma | 12.1 | 5.5 | 22.3 | 11 | ^ | ^ | ^ | ^ |
| Leukemia | 9.7 | 4.6 | 17.9 | 13 | 7.9 | 4.0 | 13.8 | 13 |
| Lymphocytic Leukemia | 4.5 | 1.4 | 10.8 | 7 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 2.8 | 1.0 | 7.3 | 6 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 5.2 | 1.7 | 12.0 | 6 | 4.9 | 2.0 | 10.0 | 8 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 3.8 | 1.3 | 8.6 | 6 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Leukemia | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 3.8 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Mesothelioma | 0.0 | 0.0 | 3.8 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Miscellaneous | 15.7 | 6.7 | 29.5 | 11 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 525.6 | 477.4 | 576.9 | 664 | 457.2 | 419.3 | 497.3 | 658 |
| Oral Cavity and Pharynx | 18.4 | 11.1 | 28.5 | 27 | 11.6 | 6.0 | 19.9 | 14 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 4.4 | 1.8 | 9.5 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Tonsil | 5.2 | 1.6 | 12.0 | 8 | 0.0 | 0.0 | 2.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Digestive System | 87.9 | 68.2 | 111.1 | 106 | 77.7 | 61.3 | 96.8 | 91 |
| Esophagus | 5.5 | 0.9 | 15.2 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 6.0 | 1.8 | 13.6 | 6 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 51.9 | 37.6 | 69.4 | 65 | 47.3 | 34.9 | 62.2 | 56 |
| Colon excluding Rectum | 33.8 | 22.3 | 48.6 | 41 | 34.8 | 24.0 | 48.4 | 38 |
| Cecum | 5.7 | 1.0 | 15.5 | 6 | 7.6 | 2.9 | 15.2 | 7 |
| Appendix | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 7.0 | 2.6 | 14.3 | 7 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 9.9 | 4.8 | 17.8 | 14 | 8.3 | 3.7 | 15.6 | 10 |
| Large Intestine, NOS | 7.0 | 2.5 | 14.9 | 8 | 4.0 | 1.3 | 9.3 | 6 |
| Rectum and Rectosigmoid Junction | 18.1 | 10.4 | 29.0 | 24 | 12.4 | 7.1 | 20.1 | 18 |
| Rectosigmoid Junction | 5.1 | 1.5 | 12.1 | 7 | ^ | ^ | ^ | ^ |
| Rectum | 13.0 | 6.7 | 22.5 | 17 | 8.8 | 4.6 | 15.1 | 14 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 7.9 | 3.0 | 16.1 | 8 | 5.6 | 2.0 | 11.8 | 7 |
| Liver | 7.4 | 2.7 | 15.6 | 7 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 948.7 | 523.3 | 1,533.8 | 29 | 383.3 | 193.7 | 661.9 | 17 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Lip | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Tongue | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Salivary Gland | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Nasopharynx | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Tonsil | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Small Intestine | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Appendix | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Ascending Colon | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Transverse Colon | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Rectum | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Liver | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 530.8 | 429.8 | 646.1 | 120 | 451.3 | 372.9 | 540.1 | 128 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Tongue | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Tonsil | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Digestive System | 162.6 | 105.7 | 235.5 | 34 | 92.8 | 59.2 | 137.2 | 26 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 27.6 | 12.4 | 54.1 | 9 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Colon and Rectum | 114.1 | 65.3 | 180.1 | 21 | 64.3 | 37.0 | 102.6 | 18 |
| Colon excluding Rectum | 96.4 | 51.4 | 159.4 | 17 | 42.9 | 21.4 | 75.4 | 12 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Sigmoid Colon | 47.5 | 19.7 | 91.6 | 10 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 21.4 | 7.3 | 47.1 | 6 |
| Rectosigmoid Junction | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Liver | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Gallbladder | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 493.5 | 301.8 | 749.0 | 25 | 300.8 | 158.0 | 504.2 | 22 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Lip | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Tongue | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Nasopharynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Tonsil | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Digestive System | 199.8 | 95.1 | 362.6 | 11 | 102.9 | 33.0 | 230.4 | 6 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Ascending Colon | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Rectum | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | 12.5 | 5.3 | 23.8 | 13 | 13.1 | 6.9 | 22.2 | 15 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 100.0 | 79.6 | 123.6 | 122 | 72.2 | 57.2 | 89.7 | 89 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Larynx | 12.5 | 6.4 | 21.6 | 15 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 87.6 | 68.4 | 109.9 | 107 | 69.8 | 55.0 | 87.1 | 85 |
| Pleura | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 3.4 | 1.4 | 7.9 | 8 | 2.6 | 0.9 | 6.3 | 6 |
| Skin excluding Basal and Squamous | 16.4 | 9.7 | 25.9 | 26 | 12.5 | 7.5 | 19.6 | 23 |
| Melanoma of the Skin | 16.1 | 9.4 | 25.6 | 25 | 12.5 | 7.5 | 19.6 | 23 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Breast | ^ | ^ | ^ | ^ | 132.2 | 113.2 | 153.4 | 210 |
| Female Genital System | 0.0 | 0.0 | 3.7 | 0 | 47.9 | 37.2 | 60.8 | 81 |
| Cervix Uteri | 0.0 | 0.0 | 3.7 | 0 | 11.3 | 7.0 | 17.5 | 23 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 3.7 | 0 | 21.5 | 14.2 | 31.1 | 33 |
| Corpus Uteri | 0.0 | 0.0 | 3.7 | 0 | 18.2 | 11.6 | 26.9 | 28 |
| Uterus, NOS | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 3.7 | 0 | 13.0 | 7.7 | 20.4 | 22 |
| Vagina | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Male Genital System | 165.1 | 137.4 | 196.2 | 191 | 0.0 | 0.0 | 2.7 | 0 |
| Prostate | 159.7 | 132.2 | 190.7 | 175 | 0.0 | 0.0 | 2.7 | 0 |
| Testis | 5.4 | 3.1 | 9.9 | 16 | 0.0 | 0.0 | 2.7 | 0 |
| Penis | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Urinary System | 56.3 | 41.1 | 74.7 | 68 | 17.5 | 10.5 | 27.1 | 22 |
| Urinary Bladder | 35.4 | 22.8 | 51.7 | 39 | 9.9 | 4.9 | 17.5 | 12 |
| Kidney and Renal Pelvis | 17.4 | 10.7 | 26.8 | 25 | 6.7 | 2.7 | 13.3 | 9 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Pancreas | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Respiratory System | 178.5 | 60.6 | 416.3 | 9 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Larynx | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Lung and Bronchus | 160.0 | 48.4 | 395.7 | 8 | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Breast | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Cervix Uteri | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Ovary | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Vagina | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Vulva | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Male Genital System | 392.4 | 175.1 | 739.9 | 11 | 0.0 | 0.0 | 92.8 | 0 |
| Prostate | 392.4 | 175.1 | 739.9 | 11 | 0.0 | 0.0 | 92.8 | 0 |
| Testis | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Penis | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 92.8 | 0 |
| Ureter | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Respiratory System | 134.2 | 88.6 | 193.2 | 32 | 70.3 | 40.5 | 111.5 | 18 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 119.4 | 76.1 | 176.3 | 28 | 66.0 | 37.3 | 106.2 | 17 |
| Pleura | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Bones and Joints | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Breast | 0.0 | 0.0 | 15.1 | 0 | 164.1 | 118.2 | 220.6 | 47 |
| Female Genital System | 0.0 | 0.0 | 15.1 | 0 | 37.6 | 18.9 | 66.6 | 12 |
| Cervix Uteri | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Ovary | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Vulva | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Male Genital System | 113.7 | 69.8 | 172.3 | 24 | 0.0 | 0.0 | 11.9 | 0 |
| Prostate | 106.1 | 63.1 | 164.1 | 21 | 0.0 | 0.0 | 11.9 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Penis | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Urinary System | 51.9 | 23.4 | 95.8 | 10 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Larynx | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Breast | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Ovary | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Vagina | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Vulva | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Male Genital System | 151.3 | 49.8 | 327.5 | 7 | 0.0 | 0.0 | 61.3 | 0 |
| Prostate | 151.3 | 49.8 | 327.5 | 7 | 0.0 | 0.0 | 61.3 | 0 |
| Testis | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Penis | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Urinary System | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Ureter | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 6.6 | 3.4 | 12.2 | 14 | 4.3 | 1.9 | 8.7 | 9 |
| Brain | 6.6 | 3.4 | 12.2 | 14 | 4.3 | 1.9 | 8.7 | 9 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 17.2 | 11.4 | 25.0 | 33 |
| Thyroid | ^ | ^ | ^ | ^ | 16.7 | 11.0 | 24.5 | 32 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 23.6 | 15.9 | 33.8 | 41 | 24.2 | 16.0 | 34.8 | 36 |
| Hodgkin Lymphoma | 5.6 | 2.7 | 11.0 | 12 | 4.9 | 2.5 | 9.2 | 12 |
| Hodgkin - Nodal | 5.6 | 2.7 | 11.0 | 12 | 4.4 | 2.2 | 8.5 | 11 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Non-Hodgkin Lymphoma | 18.0 | 11.1 | 27.5 | 29 | 19.3 | 11.7 | 29.5 | 24 |
| NHL - Nodal | 13.2 | 7.8 | 21.2 | 23 | 13.5 | 7.5 | 21.9 | 18 |
| NHL - Extranodal | 4.7 | 1.3 | 11.6 | 6 | 5.8 | 1.9 | 12.8 | 6 |
| Myeloma | ^ | ^ | ^ | ^ | 8.1 | 3.9 | 14.8 | 11 |
| Leukemia | 24.4 | 15.3 | 36.7 | 31 | 15.4 | 8.6 | 24.9 | 18 |
| Lymphocytic Leukemia | 14.5 | 7.9 | 24.1 | 19 | 5.8 | 2.1 | 12.3 | 7 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 11.1 | 5.1 | 20.5 | 12 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Myeloid and Monocytic Leukemia | 6.2 | 2.2 | 13.5 | 8 | 8.8 | 3.9 | 16.6 | 10 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 3.7 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.7 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | 8.9 | 3.7 | 17.4 | 9 | 0.0 | 0.0 | 2.7 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 3.7 | 0 | 0.0 | 0.0 | 2.7 | 0 |
| Miscellaneous | 3.1 | 1.0 | 8.0 | 6 | 12.1 | 6.0 | 21.0 | 12 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Brain | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Endocrine System | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Thyroid | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Leukemia | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 163.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |
| Miscellaneous | 0.0 | 0.0 | 163.2 | 0 | 0.0 | 0.0 | 92.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Leukemia | 23.6 | 6.2 | 57.0 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 15.1 | 0 | 0.0 | 0.0 | 11.9 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.9 | 0 |
| Miscellaneous | 0.0 | 0.0 | 15.1 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Lymphoma | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 67.9 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 67.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 595.9 | 545.3 | 649.6 | 761 | 435.5 | 399.4 | 473.7 | 643 |
| Oral Cavity and Pharynx | 13.5 | 7.9 | 21.5 | 23 | 7.8 | 3.7 | 14.0 | 12 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Digestive System | 91.7 | 72.9 | 113.5 | 118 | 57.6 | 44.4 | 73.1 | 76 |
| Esophagus | 9.8 | 4.3 | 18.4 | 12 | ^ | ^ | ^ | ^ |
| Stomach | 7.0 | 2.7 | 14.5 | 8 | 4.4 | 1.4 | 9.9 | 6 |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 46.8 | 34.2 | 62.1 | 61 | 31.6 | 21.9 | 43.7 | 41 |
| Colon excluding Rectum | 32.3 | 21.8 | 45.8 | 42 | 24.5 | 16.0 | 35.5 | 31 |
| Cecum | 4.7 | 1.6 | 10.8 | 7 | 7.6 | 3.1 | 15.0 | 8 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 6.8 | 2.5 | 14.3 | 8 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 11.6 | 5.3 | 21.2 | 12 | 6.2 | 2.5 | 12.6 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 14.4 | 8.3 | 23.3 | 19 | 7.1 | 3.2 | 13.4 | 10 |
| Rectosigmoid Junction | 5.1 | 1.8 | 11.5 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 9.3 | 4.6 | 16.7 | 13 | 6.7 | 2.9 | 13.0 | 9 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 5.6 | 2.2 | 11.5 | 8 |
| Liver and Intrahepatic Bile Duct | 9.7 | 3.9 | 19.0 | 14 | ^ | ^ | ^ | ^ |
| Liver | 5.6 | 2.6 | 10.9 | 11 | 0.0 | 0.0 | 2.6 | 0 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 653.1 | 403.5 | 997.4 | 28 | 462.0 | 260.4 | 743.2 | 21 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Tongue | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Nasopharynx | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Tonsil | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Small Intestine | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Appendix | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Ascending Colon | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Transverse Colon | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Rectum | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Liver | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 654.6 | 542.9 | 780.1 | 153 | 459.5 | 381.7 | 547.5 | 136 |
| Oral Cavity and Pharynx | 31.8 | 11.5 | 66.9 | 7 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Salivary Gland | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Oropharynx | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Hypopharynx | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Digestive System | 277.1 | 205.4 | 363.0 | 62 | 172.1 | 123.7 | 231.6 | 46 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 28.5 | 12.8 | 55.7 | 9 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 166.6 | 112.6 | 234.9 | 37 | 93.2 | 58.8 | 138.6 | 25 |
| Colon excluding Rectum | 120.8 | 73.7 | 183.3 | 24 | 58.6 | 31.5 | 97.4 | 15 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 52.0 | 21.8 | 99.2 | 9 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 45.8 | 23.2 | 80.6 | 13 | 34.5 | 16.1 | 63.5 | 10 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 36.1 | 16.2 | 68.6 | 10 | 28.3 | 11.8 | 55.7 | 8 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 524.8 | 339.9 | 764.5 | 33 | 259.1 | 172.5 | 378.3 | 31 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Lip | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Tongue | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Salivary Gland | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Oropharynx | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Digestive System | 81.6 | 31.6 | 176.6 | 7 | 69.6 | 25.9 | 153.4 | 7 |
| Esophagus | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Stomach | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Descending Colon | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Gallbladder | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Pancreas | 13.5 | 6.4 | 24.0 | 16 | 8.0 | 3.6 | 14.8 | 10 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 107.8 | 86.9 | 131.6 | 130 | 56.6 | 43.6 | 71.9 | 73 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 11.9 | 5.4 | 21.9 | 15 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 95.9 | 76.4 | 118.3 | 115 | 55.6 | 42.7 | 70.9 | 71 |
| Pleura | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 19.7 | 11.4 | 31.0 | 32 | 13.7 | 9.0 | 20.2 | 29 |
| Melanoma of the Skin | 18.8 | 10.7 | 30.1 | 30 | 13.3 | 8.6 | 19.7 | 28 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 148.6 | 128.5 | 170.8 | 233 |
| Female Genital System | 0.0 | 0.0 | 3.5 | 0 | 57.7 | 45.0 | 72.7 | 84 |
| Cervix Uteri | 0.0 | 0.0 | 3.5 | 0 | 10.5 | 6.4 | 16.4 | 22 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 3.5 | 0 | 25.8 | 17.1 | 36.8 | 33 |
| Corpus Uteri | 0.0 | 0.0 | 3.5 | 0 | 23.2 | 15.1 | 33.7 | 30 |
| Uterus, NOS | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 3.5 | 0 | 17.9 | 10.9 | 27.3 | 23 |
| Vagina | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 190.0 | 161.8 | 221.1 | 222 | 0.0 | 0.0 | 2.6 | 0 |
| Prostate | 183.7 | 155.7 | 214.7 | 205 | 0.0 | 0.0 | 2.6 | 0 |
| Testis | 6.3 | 3.7 | 11.0 | 17 | 0.0 | 0.0 | 2.6 | 0 |
| Penis | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Urinary System | 85.7 | 64.9 | 110.0 | 91 | 25.8 | 17.7 | 36.1 | 37 |
| Urinary Bladder | 68.2 | 48.9 | 91.4 | 64 | 14.6 | 8.5 | 23.2 | 19 |
| Kidney and Renal Pelvis | 15.2 | 9.3 | 23.5 | 25 | 11.1 | 6.3 | 18.1 | 18 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Pancreas | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Larynx | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Breast | 0.0 | 0.0 | 139.2 | 0 | 133.3 | 37.4 | 315.6 | 6 |
| Female Genital System | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Ovary | 0.0 | 0.0 | 139.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Vulva | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Male Genital System | 272.6 | 132.7 | 506.2 | 12 | 0.0 | 0.0 | 85.8 | 0 |
| Prostate | 266.9 | 127.8 | 500.5 | 11 | 0.0 | 0.0 | 85.8 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Penis | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | 31.1 | 12.1 | 62.5 | 7 |
| Retroperitoneum | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Respiratory System | 159.6 | 107.5 | 225.8 | 37 | 68.3 | 39.6 | 108.1 | 18 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Larynx | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 147.2 | 97.4 | 211.0 | 34 | 64.2 | 36.5 | 103.0 | 17 |
| Pleura | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Breast | 0.0 | 0.0 | 14.9 | 0 | 118.7 | 83.0 | 163.9 | 38 |
| Female Genital System | 0.0 | 0.0 | 14.9 | 0 | 27.7 | 12.4 | 52.9 | 9 |
| Cervix Uteri | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Ovary | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Vulva | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Male Genital System | 88.8 | 48.0 | 145.8 | 17 | 0.0 | 0.0 | 11.5 | 0 |
| Prostate | 86.8 | 46.2 | 143.8 | 16 | 0.0 | 0.0 | 11.5 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Penis | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Urinary System | 46.6 | 20.9 | 86.6 | 11 | 26.3 | 9.2 | 55.9 | 6 |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 34.0 | 11.8 | 72.1 | 7 | ^ | ^ | ^ | ^ |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 98.6 | 39.9 | 203.8 | 8 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Lung and Bronchus | 98.6 | 39.9 | 203.8 | 8 | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Bones and Joints | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Breast | 0.0 | 0.0 | 59.2 | 0 | 73.1 | 34.4 | 146.4 | 10 |
| Female Genital System | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Ovary | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Vulva | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 214.5 | 89.3 | 409.5 | 9 | 0.0 | 0.0 | 52.1 | 0 |
| Prostate | 208.5 | 84.4 | 403.5 | 8 | 0.0 | 0.0 | 52.1 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Penis | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Ureter | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 6.1 | 2.5 | 12.4 | 12 | 7.5 | 3.8 | 13.3 | 13 |
| Brain | 6.1 | 2.5 | 12.4 | 12 | 7.5 | 3.8 | 13.3 | 13 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Endocrine System | 3.8 | 1.8 | 8.2 | 10 | 11.0 | 6.6 | 17.4 | 23 |
| Thyroid | 3.5 | 1.6 | 7.8 | 9 | 11.0 | 6.6 | 17.4 | 23 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Lymphoma | 36.8 | 26.7 | 49.3 | 63 | 19.2 | 11.7 | 29.3 | 24 |
| Hodgkin Lymphoma | 4.1 | 1.8 | 8.8 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 4.1 | 1.8 | 8.8 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Non-Hodgkin Lymphoma | 32.7 | 23.1 | 44.9 | 54 | 17.6 | 10.4 | 27.5 | 21 |
| NHL - Nodal | 22.6 | 14.7 | 33.1 | 37 | 12.3 | 6.3 | 20.9 | 15 |
| NHL - Extranodal | 10.1 | 5.2 | 17.7 | 17 | 5.4 | 1.9 | 11.5 | 6 |
| Myeloma | ^ | ^ | ^ | ^ | 3.4 | 1.2 | 7.6 | 6 |
| Leukemia | 18.2 | 10.5 | 29.1 | 28 | 12.7 | 6.8 | 21.1 | 16 |
| Lymphocytic Leukemia | 11.4 | 5.0 | 21.5 | 15 | 5.7 | 2.0 | 12.0 | 7 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 9.4 | 3.4 | 19.4 | 10 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 5.3 | 2.2 | 10.9 | 10 | 4.4 | 1.4 | 10.0 | 6 |
| Acute Myeloid Leukemia | 3.8 | 1.2 | 9.2 | 6 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 3.5 | 0 | 0.0 | 0.0 | 2.6 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 3.5 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.6 | 0 |
| Miscellaneous | 9.1 | 3.3 | 18.7 | 11 | 9.0 | 4.2 | 16.4 | 10 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Brain | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Thyroid | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 85.8 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Myeloma | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |
| Miscellaneous | 0.0 | 0.0 | 139.2 | 0 | 0.0 | 0.0 | 85.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Lymphoma | 16.4 | 5.9 | 38.1 | 6 | 14.1 | 4.9 | 32.6 | 6 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 14.1 | 4.9 | 32.6 | 6 |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 14.9 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Mesothelioma | 0.0 | 0.0 | 14.9 | 0 | 0.0 | 0.0 | 11.5 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.5 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Brain | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 52.1 | 0 |
| Leukemia | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Mesothelioma | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |
| Miscellaneous | 0.0 | 0.0 | 59.2 | 0 | 0.0 | 0.0 | 52.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 575.1 | 527.8 | 625.0 | 766 | 477.0 | 440.4 | 515.7 | 739 |
| Oral Cavity and Pharynx | 9.1 | 5.3 | 15.0 | 20 | 8.9 | 4.6 | 15.5 | 14 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 2.9 | 1.0 | 7.1 | 6 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Hypopharynx | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Digestive System | 104.7 | 84.1 | 128.3 | 132 | 75.7 | 60.5 | 93.1 | 98 |
| Esophagus | 5.0 | 2.0 | 10.3 | 8 | ^ | ^ | ^ | ^ |
| Stomach | 15.6 | 7.3 | 27.8 | 13 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 53.8 | 39.1 | 71.5 | 66 | 49.3 | 37.2 | 63.9 | 62 |
| Colon excluding Rectum | 39.0 | 25.8 | 55.7 | 41 | 33.9 | 23.8 | 46.4 | 40 |
| Cecum | 8.9 | 2.7 | 19.5 | 8 | 13.9 | 7.8 | 22.6 | 16 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Ascending Colon | 11.1 | 4.0 | 22.6 | 8 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 7.9 | 3.6 | 14.8 | 10 | 10.7 | 5.3 | 18.6 | 12 |
| Large Intestine, NOS | 4.0 | 1.2 | 9.4 | 6 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 14.8 | 9.3 | 22.5 | 25 | 15.5 | 9.3 | 23.9 | 22 |
| Rectosigmoid Junction | 5.0 | 1.8 | 10.8 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 9.8 | 5.7 | 16.0 | 19 | 12.4 | 6.8 | 20.3 | 17 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 6.3 | 3.1 | 11.8 | 12 | 5.2 | 1.7 | 11.4 | 6 |
| Liver | 6.3 | 3.1 | 11.8 | 12 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 753.3 | 464.8 | 1,148.4 | 28 | 626.5 | 375.4 | 962.1 | 28 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Lip | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Salivary Gland | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Nasopharynx | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Tonsil | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Oropharynx | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Hypopharynx | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Digestive System | 260.5 | 102.7 | 537.7 | 9 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Small Intestine | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Appendix | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Ascending Colon | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Descending Colon | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Rectum | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 504.0 | 409.6 | 611.6 | 129 | 535.8 | 452.7 | 628.7 | 166 |
| Oral Cavity and Pharynx | 46.1 | 21.9 | 83.6 | 13 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Digestive System | 171.1 | 117.8 | 238.0 | 43 | 208.4 | 155.0 | 272.7 | 56 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 23.6 | 9.7 | 48.8 | 8 | 17.1 | 5.7 | 38.5 | 6 |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 96.7 | 57.0 | 150.7 | 22 | 141.0 | 96.6 | 196.6 | 36 |
| Colon excluding Rectum | 78.8 | 42.4 | 130.3 | 17 | 109.1 | 70.4 | 159.3 | 27 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Sigmoid Colon | 29.9 | 10.4 | 64.0 | 7 | 39.4 | 18.8 | 70.9 | 11 |
| Large Intestine, NOS | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 31.8 | 13.5 | 61.4 | 9 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 20.1 | 6.5 | 44.8 | 6 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 494.1 | 302.5 | 746.8 | 27 | 202.1 | 125.6 | 311.2 | 23 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Lip | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Tongue | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Salivary Gland | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Nasopharynx | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Tonsil | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Oropharynx | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hypopharynx | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Digestive System | 79.9 | 30.3 | 174.4 | 7 | 53.8 | 19.2 | 124.8 | 6 |
| Esophagus | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Cecum | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Appendix | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Ascending Colon | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Transverse Colon | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Descending Colon | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Rectum | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Gallbladder | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 19.9 | 11.9 | 30.8 | 26 | 8.2 | 4.1 | 14.4 | 13 |
| Retroperitoneum | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 96.5 | 78.1 | 117.5 | 127 | 79.6 | 64.4 | 97.1 | 104 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 4.1 | 1.6 | 9.2 | 8 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 91.6 | 73.6 | 112.3 | 118 | 77.7 | 62.7 | 95.1 | 101 |
| Pleura | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 3.9 | 1.6 | 8.2 | 8 |
| Skin excluding Basal and Squamous | 16.9 | 11.0 | 25.0 | 31 | 9.5 | 5.7 | 15.0 | 21 |
| Melanoma of the Skin | 14.6 | 9.2 | 22.3 | 27 | 6.6 | 3.7 | 11.1 | 16 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 156.7 | 136.5 | 178.9 | 261 |
| Female Genital System | 0.0 | 0.0 | 3.2 | 0 | 52.7 | 41.3 | 66.2 | 87 |
| Cervix Uteri | 0.0 | 0.0 | 3.2 | 0 | 13.2 | 8.0 | 20.3 | 25 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 3.2 | 0 | 25.7 | 17.9 | 35.5 | 40 |
| Corpus Uteri | 0.0 | 0.0 | 3.2 | 0 | 25.7 | 17.9 | 35.5 | 40 |
| Uterus, NOS | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Ovary | 0.0 | 0.0 | 3.2 | 0 | 9.7 | 5.3 | 16.3 | 17 |
| Vagina | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 186.8 | 159.9 | 216.5 | 232 | 0.0 | 0.0 | 2.4 | 0 |
| Prostate | 178.2 | 152.1 | 207.2 | 216 | 0.0 | 0.0 | 2.4 | 0 |
| Testis | 3.9 | 1.9 | 7.9 | 11 | 0.0 | 0.0 | 2.4 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Urinary System | 62.9 | 47.7 | 80.8 | 77 | 16.5 | 10.3 | 24.8 | 25 |
| Urinary Bladder | 46.4 | 32.9 | 63.0 | 49 | 10.2 | 5.3 | 17.3 | 14 |
| Kidney and Renal Pelvis | 14.6 | 9.0 | 22.5 | 26 | 6.3 | 3.0 | 11.6 | 11 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Bones and Joints | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Breast | ^ | ^ | ^ | ^ | 259.1 | 119.4 | 476.0 | 15 |
| Female Genital System | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Ovary | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Vulva | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Male Genital System | 222.2 | 88.2 | 468.5 | 9 | 0.0 | 0.0 | 84.5 | 0 |
| Prostate | 222.2 | 88.2 | 468.5 | 9 | 0.0 | 0.0 | 84.5 | 0 |
| Testis | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Penis | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 157.9 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Ureter | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Respiratory System | 131.0 | 87.5 | 187.1 | 34 | 69.0 | 40.5 | 108.2 | 19 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Lung and Bronchus | 124.9 | 82.2 | 180.3 | 32 | 65.9 | 38.0 | 104.5 | 18 |
| Pleura | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Bones and Joints | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Breast | 0.0 | 0.0 | 14.6 | 0 | 162.2 | 118.9 | 215.2 | 51 |
| Female Genital System | 0.0 | 0.0 | 14.6 | 0 | 34.2 | 18.4 | 58.5 | 14 |
| Cervix Uteri | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 14.6 | 0 | 15.5 | 5.6 | 34.7 | 6 |
| Corpus Uteri | 0.0 | 0.0 | 14.6 | 0 | 15.5 | 5.6 | 34.7 | 6 |
| Uterus, NOS | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Ovary | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Vulva | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Male Genital System | 75.8 | 40.2 | 126.4 | 16 | 0.0 | 0.0 | 11.1 | 0 |
| Prostate | 62.6 | 31.8 | 107.9 | 14 | 0.0 | 0.0 | 11.1 | 0 |
| Testis | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Urinary System | 36.3 | 15.2 | 70.8 | 10 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Kidney and Renal Pelvis | 27.6 | 9.2 | 60.5 | 7 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Pancreas | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Bones and Joints | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Breast | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Ovary | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Vulva | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Male Genital System | 167.8 | 59.9 | 345.9 | 7 | 0.0 | 0.0 | 48.9 | 0 |
| Prostate | 167.8 | 59.9 | 345.9 | 7 | 0.0 | 0.0 | 48.9 | 0 |
| Testis | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Penis | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Brain and Other Nervous System | 9.0 | 5.1 | 15.0 | 19 | 6.4 | 3.1 | 11.7 | 12 |
| Brain | 9.0 | 5.1 | 15.0 | 19 | 6.4 | 3.1 | 11.7 | 12 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Endocrine System | 3.1 | 1.3 | 7.1 | 8 | 16.7 | 11.4 | 23.7 | 35 |
| Thyroid | 3.1 | 1.3 | 7.1 | 8 | 16.2 | 11.0 | 23.2 | 34 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 3.2 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | 32.6 | 22.0 | 46.0 | 48 | 23.5 | 16.1 | 33.0 | 38 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 3.2 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Non-Hodgkin Lymphoma | 29.2 | 19.2 | 42.1 | 44 | 21.9 | 14.8 | 31.1 | 35 |
| NHL - Nodal | 20.2 | 12.4 | 30.7 | 32 | 10.5 | 5.9 | 17.2 | 17 |
| NHL - Extranodal | 9.1 | 3.6 | 17.8 | 12 | 11.4 | 6.4 | 18.5 | 18 |
| Myeloma | 9.9 | 5.3 | 17.0 | 15 | 6.7 | 2.9 | 12.7 | 9 |
| Leukemia | 26.6 | 16.3 | 40.3 | 31 | 8.5 | 4.1 | 15.2 | 12 |
| Lymphocytic Leukemia | 13.3 | 6.3 | 23.6 | 15 | 5.2 | 1.9 | 10.8 | 7 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 8.0 | 3.2 | 15.9 | 8 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 11.3 | 4.9 | 21.2 | 13 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 6.2 | 2.1 | 13.3 | 7 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Mesothelioma | 6.2 | 2.1 | 13.4 | 6 | 0.0 | 0.0 | 2.4 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Miscellaneous | 5.9 | 2.5 | 11.8 | 9 | 11.3 | 6.0 | 19.0 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Brain | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Endocrine System | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Thyroid | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 84.5 | 0 |
| Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Mesothelioma | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |
| Miscellaneous | 0.0 | 0.0 | 157.9 | 0 | 0.0 | 0.0 | 84.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 14.6 | 0 | 0.0 | 0.0 | 11.1 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 11.1 | 0 |
| Miscellaneous | 0.0 | 0.0 | 14.6 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 58.3 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Mesothelioma | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Miscellaneous | 0.0 | 0.0 | 58.3 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 559.0 | 514.8 | 605.7 | 811 | 455.7 | 420.1 | 493.3 | 710 |
| Oral Cavity and Pharynx | 14.0 | 8.0 | 22.3 | 26 | 7.0 | 2.9 | 13.6 | 8 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Tongue | 6.9 | 2.5 | 14.4 | 11 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 3.0 | 0 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.0 | 0.0 | 3.0 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Digestive System | 97.4 | 79.2 | 118.2 | 137 | 67.1 | 52.7 | 83.8 | 84 |
| Esophagus | 10.0 | 5.4 | 16.9 | 16 | ^ | ^ | ^ | ^ |
| Stomach | 8.6 | 3.0 | 18.1 | 10 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 59.6 | 45.5 | 76.3 | 79 | 39.1 | 28.2 | 52.4 | 47 |
| Colon excluding Rectum | 49.0 | 36.0 | 64.7 | 60 | 29.6 | 20.2 | 41.5 | 35 |
| Cecum | 8.0 | 3.6 | 15.0 | 12 | 11.5 | 6.0 | 19.5 | 13 |
| Appendix | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Ascending Colon | 14.1 | 7.7 | 23.1 | 16 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 4.5 | 1.5 | 10.0 | 7 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 13.0 | 6.5 | 22.5 | 14 | 5.4 | 1.9 | 11.5 | 6 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 10.6 | 5.9 | 17.6 | 19 | 9.5 | 4.6 | 16.7 | 12 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 7.9 | 3.8 | 14.5 | 14 | 7.7 | 3.4 | 14.4 | 10 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 3.0 | 0 | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 8.1 | 3.6 | 15.3 | 12 | ^ | ^ | ^ | ^ |
| Liver | 8.1 | 3.6 | 15.3 | 12 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 732.3 | 479.3 | 1,071.9 | 31 | 409.7 | 227.6 | 664.8 | 22 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Lip | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Tongue | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Salivary Gland | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Nasopharynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Tonsil | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Oropharynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Hypopharynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Appendix | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Transverse Colon | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Descending Colon | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Rectum | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Gallbladder | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 622.2 | 517.5 | 739.8 | 155 | 505.9 | 427.4 | 593.6 | 165 |
| Oral Cavity and Pharynx | 48.1 | 24.3 | 84.0 | 14 | ^ | ^ | ^ | ^ |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Nasopharynx | 18.2 | 5.9 | 42.4 | 6 | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Oropharynx | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Hypopharynx | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Digestive System | 253.1 | 185.5 | 334.7 | 59 | 194.8 | 146.3 | 253.0 | 59 |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Stomach | 25.4 | 9.6 | 53.3 | 7 | 33.2 | 15.6 | 60.7 | 11 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Colon and Rectum | 157.7 | 106.8 | 222.0 | 38 | 126.2 | 88.8 | 173.2 | 39 |
| Colon excluding Rectum | 114.7 | 73.3 | 168.9 | 29 | 93.3 | 61.8 | 134.2 | 29 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | 21.9 | 7.6 | 47.0 | 6 |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 55.5 | 30.8 | 91.6 | 16 | 25.7 | 11.7 | 48.7 | 9 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 43.1 | 17.5 | 84.1 | 9 | 32.9 | 15.2 | 60.7 | 10 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 37.0 | 13.1 | 77.3 | 7 | 25.0 | 9.6 | 51.1 | 7 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 336.1 | 203.5 | 515.2 | 22 | 297.8 | 175.5 | 463.4 | 30 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Nasopharynx | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Digestive System | 110.7 | 42.2 | 227.3 | 8 | 86.6 | 23.3 | 203.0 | 7 |
| Esophagus | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 43.8 | 0 |
| Cecum | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Appendix | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Ascending Colon | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Transverse Colon | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 43.8 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Rectum | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 8.2 | 3.9 | 14.9 | 13 | 16.0 | 9.5 | 24.9 | 20 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 89.8 | 73.1 | 108.8 | 127 | 82.3 | 67.0 | 99.8 | 111 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Larynx | 5.8 | 2.5 | 11.5 | 9 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 82.8 | 66.7 | 101.4 | 115 | 79.7 | 64.6 | 97.0 | 107 |
| Pleura | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 22.8 | 14.1 | 34.3 | 35 | 11.6 | 6.9 | 18.2 | 22 |
| Melanoma of the Skin | 19.3 | 11.2 | 30.4 | 29 | 9.3 | 5.4 | 15.0 | 19 |
| Other Non-Epithelial Skin | 3.4 | 1.0 | 8.4 | 6 | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 146.6 | 127.7 | 167.3 | 254 |
| Female Genital System | 0.0 | 0.0 | 3.0 | 0 | 51.2 | 40.1 | 64.2 | 85 |
| Cervix Uteri | 0.0 | 0.0 | 3.0 | 0 | 4.7 | 2.0 | 9.4 | 9 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 3.0 | 0 | 21.8 | 15.1 | 30.4 | 38 |
| Corpus Uteri | 0.0 | 0.0 | 3.0 | 0 | 21.8 | 15.1 | 30.4 | 38 |
| Uterus, NOS | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Ovary | 0.0 | 0.0 | 3.0 | 0 | 17.5 | 11.1 | 26.0 | 27 |
| Vagina | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Vulva | 0.0 | 0.0 | 3.0 | 0 | 4.6 | 1.7 | 9.9 | 7 |
| Other Female Genital Organs | 0.0 | 0.0 | 3.0 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 191.5 | 166.0 | 219.6 | 264 | 0.0 | 0.0 | 2.4 | 0 |
| Prostate | 182.1 | 156.8 | 209.9 | 239 | 0.0 | 0.0 | 2.4 | 0 |
| Testis | 9.5 | 6.1 | 14.6 | 25 | 0.0 | 0.0 | 2.4 | 0 |
| Penis | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Urinary System | 61.4 | 46.7 | 78.8 | 83 | 15.8 | 9.6 | 24.3 | 22 |
| Urinary Bladder | 40.1 | 27.8 | 55.4 | 47 | 9.2 | 4.4 | 16.3 | 11 |
| Kidney and Renal Pelvis | 20.4 | 13.2 | 29.9 | 35 | 6.6 | 3.1 | 12.2 | 11 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Larynx | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Bones and Joints | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Breast | 0.0 | 0.0 | 139.1 | 0 | 157.6 | 61.3 | 321.9 | 9 |
| Female Genital System | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Ovary | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Vagina | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Vulva | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Male Genital System | 361.1 | 197.8 | 612.6 | 16 | 0.0 | 0.0 | 75.6 | 0 |
| Prostate | 361.1 | 197.8 | 612.6 | 16 | 0.0 | 0.0 | 75.6 | 0 |
| Testis | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Penis | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Pancreas | 31.4 | 9.4 | 70.5 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Respiratory System | 107.3 | 67.9 | 159.3 | 26 | 62.7 | 36.9 | 98.4 | 19 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 99.9 | 61.9 | 150.8 | 24 | 51.3 | 28.0 | 84.8 | 15 |
| Pleura | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Bones and Joints | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Breast | 0.0 | 0.0 | 14.3 | 0 | 87.4 | 58.8 | 124.6 | 32 |
| Female Genital System | 0.0 | 0.0 | 14.3 | 0 | 41.8 | 22.6 | 70.2 | 16 |
| Cervix Uteri | 0.0 | 0.0 | 14.3 | 0 | 18.5 | 6.7 | 39.7 | 7 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 14.3 | 0 | 14.3 | 5.1 | 32.1 | 6 |
| Corpus Uteri | 0.0 | 0.0 | 14.3 | 0 | 14.3 | 5.1 | 32.1 | 6 |
| Uterus, NOS | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Ovary | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Male Genital System | 83.3 | 49.5 | 129.5 | 21 | 0.0 | 0.0 | 10.6 | 0 |
| Prostate | 80.8 | 47.5 | 126.9 | 20 | 0.0 | 0.0 | 10.6 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Penis | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Urinary System | 67.2 | 35.2 | 113.4 | 16 | 39.8 | 19.2 | 71.0 | 11 |
| Urinary Bladder | 29.0 | 10.4 | 61.5 | 7 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 35.2 | 12.8 | 72.8 | 8 | 25.9 | 10.7 | 51.1 | 8 |
| Ureter | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | ^ | ^ | ^ | ^ | 49.8 | 17.7 | 115.3 | 6 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Breast | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Ovary | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Vagina | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Vulva | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Male Genital System | 110.2 | 42.3 | 225.7 | 7 | 0.0 | 0.0 | 43.8 | 0 |
| Prostate | 110.2 | 42.3 | 225.7 | 7 | 0.0 | 0.0 | 43.8 | 0 |
| Testis | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Penis | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Urinary System | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Ureter | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Brain and Other Nervous System | 6.2 | 3.0 | 11.6 | 13 | 8.9 | 4.8 | 15.1 | 15 |
| Brain | 6.2 | 3.0 | 11.6 | 13 | 8.9 | 4.8 | 15.1 | 15 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Endocrine System | 8.3 | 4.7 | 14.0 | 19 | 8.7 | 5.2 | 13.9 | 20 |
| Thyroid | 8.0 | 4.4 | 13.6 | 18 | 8.7 | 5.2 | 13.9 | 20 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Lymphoma | 26.8 | 18.2 | 37.7 | 47 | 24.1 | 17.1 | 33.0 | 44 |
| Hodgkin Lymphoma | 2.9 | 1.2 | 6.6 | 8 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 2.9 | 1.2 | 6.6 | 8 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Non-Hodgkin Lymphoma | 23.9 | 15.6 | 34.7 | 39 | 21.9 | 15.2 | 30.5 | 39 |
| NHL - Nodal | 18.3 | 11.7 | 27.2 | 32 | 17.7 | 11.7 | 25.7 | 31 |
| NHL - Extranodal | 5.6 | 1.6 | 13.1 | 7 | 4.2 | 1.7 | 8.5 | 8 |
| Myeloma | 9.1 | 3.9 | 17.2 | 11 | ^ | ^ | ^ | ^ |
| Leukemia | 12.1 | 6.7 | 20.1 | 20 | 16.0 | 9.5 | 24.9 | 22 |
| Lymphocytic Leukemia | 8.6 | 4.3 | 15.3 | 15 | 7.3 | 3.1 | 14.0 | 9 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 6.6 | 3.3 | 12.1 | 13 | 6.5 | 2.5 | 13.0 | 7 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 4.9 | 1.8 | 10.3 | 8 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 3.3 | 1.0 | 7.8 | 6 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 3.0 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 3.0 | 0 | ^ | ^ | ^ | ^ |
| Mesothelioma | 8.0 | 2.9 | 16.5 | 8 | 0.0 | 0.0 | 2.4 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Miscellaneous | 6.2 | 2.7 | 12.2 | 9 | 9.7 | 4.8 | 17.0 | 12 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Endocrine System | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Myeloma | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 139.1 | 0 | ^ | ^ | ^ | ^ |
| Other Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Mesothelioma | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 139.1 | 0 | 0.0 | 0.0 | 75.6 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 75.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 15.6 | 6.2 | 33.5 | 7 |
| Thyroid | ^ | ^ | ^ | ^ | 15.6 | 6.2 | 33.5 | 7 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Lymphoma | 12.6 | 4.5 | 31.2 | 6 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Non-Hodgkin Lymphoma | 12.6 | 4.5 | 31.2 | 6 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.6 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Leukemia | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Mesothelioma | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 14.3 | 0 | 0.0 | 0.0 | 10.6 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Endocrine System | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Lymphoma | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 50.9 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |
| Miscellaneous | 0.0 | 0.0 | 50.9 | 0 | 0.0 | 0.0 | 43.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 604.6 | 558.5 | 653.1 | 864 | 463.7 | 428.6 | 500.7 | 743 |
| Oral Cavity and Pharynx | 17.9 | 11.2 | 27.0 | 29 | 6.3 | 2.6 | 12.4 | 8 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 6.7 | 2.8 | 13.2 | 12 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Digestive System | 116.4 | 96.8 | 138.4 | 164 | 78.0 | 63.0 | 95.2 | 103 |
| Esophagus | 12.7 | 7.0 | 20.8 | 19 | ^ | ^ | ^ | ^ |
| Stomach | 9.8 | 4.6 | 17.6 | 12 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 68.2 | 53.3 | 85.5 | 97 | 55.6 | 42.9 | 70.6 | 71 |
| Colon excluding Rectum | 49.1 | 36.0 | 64.8 | 62 | 40.3 | 29.5 | 53.4 | 50 |
| Cecum | 14.6 | 7.6 | 24.7 | 15 | 10.9 | 5.7 | 18.5 | 13 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 7.8 | 3.4 | 14.8 | 10 | 8.7 | 4.0 | 15.8 | 10 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | 6.4 | 2.4 | 13.1 | 8 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 14.6 | 7.9 | 24.2 | 19 | 12.8 | 7.0 | 20.8 | 16 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 19.1 | 12.7 | 27.6 | 35 | 15.3 | 9.2 | 23.7 | 21 |
| Rectosigmoid Junction | 4.0 | 1.6 | 8.5 | 8 | ^ | ^ | ^ | ^ |
| Rectum | 15.1 | 9.4 | 22.9 | 27 | 12.0 | 6.7 | 19.6 | 17 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 11.7 | 6.5 | 19.2 | 18 | ^ | ^ | ^ | ^ |
| Liver | 10.3 | 5.5 | 17.5 | 16 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 726.7 | 475.8 | 1,056.2 | 35 | 236.9 | 123.9 | 406.9 | 16 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Lip | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Tongue | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Salivary Gland | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Nasopharynx | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Tonsil | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Oropharynx | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Hypopharynx | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Digestive System | 185.6 | 72.0 | 385.6 | 9 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Small Intestine | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Cecum | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Appendix | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Ascending Colon | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Transverse Colon | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Descending Colon | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Rectum | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Liver | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Gallbladder | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 487.7 | 399.3 | 588.1 | 131 | 488.6 | 413.0 | 573.2 | 162 |
| Oral Cavity and Pharynx | 25.2 | 9.8 | 52.2 | 7 | ^ | ^ | ^ | ^ |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Salivary Gland | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Oropharynx | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Digestive System | 163.4 | 114.2 | 224.8 | 44 | 164.5 | 120.0 | 218.8 | 48 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 37.5 | 14.6 | 74.9 | 8 | 19.1 | 7.0 | 40.7 | 6 |
| Small Intestine | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Colon and Rectum | 89.8 | 54.0 | 138.2 | 24 | 95.0 | 61.1 | 139.1 | 26 |
| Colon excluding Rectum | 54.0 | 27.3 | 93.3 | 14 | 66.8 | 39.0 | 105.0 | 18 |
| Cecum | ^ | ^ | ^ | ^ | 20.4 | 7.2 | 43.7 | 6 |
| Appendix | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 24.4 | 8.8 | 51.0 | 6 |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Sigmoid Colon | 27.9 | 10.1 | 58.9 | 7 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Rectum and Rectosigmoid Junction | 35.8 | 15.0 | 69.6 | 10 | 28.1 | 11.6 | 55.1 | 8 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 30.4 | 11.1 | 63.3 | 8 | 23.1 | 8.8 | 47.5 | 7 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Gallbladder | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 408.3 | 253.5 | 611.3 | 28 | 374.3 | 265.6 | 510.8 | 48 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Lip | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Tongue | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Salivary Gland | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Nasopharynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Tonsil | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Hypopharynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Digestive System | 170.7 | 80.3 | 307.5 | 12 | 118.0 | 53.1 | 218.5 | 12 |
| Esophagus | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 53.6 | 19.4 | 117.1 | 6 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Ascending Colon | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Transverse Colon | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Descending Colon | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 6.6 | 2.6 | 13.2 | 9 | 7.9 | 3.8 | 14.1 | 11 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Respiratory System | 94.1 | 76.2 | 114.6 | 125 | 55.2 | 43.4 | 69.1 | 82 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Larynx | 4.2 | 1.7 | 9.0 | 8 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 88.3 | 70.8 | 108.4 | 115 | 53.4 | 41.8 | 67.1 | 79 |
| Pleura | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.3 | 1.4 | 9.6 | 8 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 13.9 | 8.2 | 22.0 | 29 | 12.7 | 7.6 | 19.7 | 23 |
| Melanoma of the Skin | 13.9 | 8.2 | 22.0 | 29 | 11.6 | 6.9 | 18.3 | 22 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 157.9 | 138.7 | 179.1 | 277 |
| Female Genital System | 0.0 | 0.0 | 2.8 | 0 | 51.5 | 40.9 | 64.0 | 93 |
| Cervix Uteri | 0.0 | 0.0 | 2.8 | 0 | 6.4 | 3.6 | 11.0 | 15 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.8 | 0 | 24.8 | 17.6 | 33.8 | 45 |
| Corpus Uteri | 0.0 | 0.0 | 2.8 | 0 | 24.4 | 17.3 | 33.4 | 44 |
| Uterus, NOS | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 2.8 | 0 | 15.6 | 9.9 | 23.4 | 26 |
| Vagina | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 203.4 | 177.1 | 232.2 | 277 | 0.0 | 0.0 | 2.3 | 0 |
| Prostate | 197.5 | 171.4 | 226.1 | 263 | 0.0 | 0.0 | 2.3 | 0 |
| Testis | 4.4 | 2.3 | 8.4 | 12 | 0.0 | 0.0 | 2.3 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Urinary System | 60.4 | 45.1 | 78.6 | 78 | 24.1 | 16.0 | 34.6 | 30 |
| Urinary Bladder | 46.6 | 32.8 | 63.4 | 55 | 14.0 | 7.8 | 22.5 | 16 |
| Kidney and Renal Pelvis | 13.8 | 7.9 | 22.2 | 23 | 10.2 | 5.3 | 17.2 | 14 |
| Ureter | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Respiratory System | 198.2 | 72.3 | 418.5 | 7 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Lung and Bronchus | 153.6 | 50.6 | 347.4 | 6 | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Bones and Joints | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Breast | 0.0 | 0.0 | 110.2 | 0 | 119.7 | 40.2 | 264.2 | 7 |
| Female Genital System | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Ovary | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Male Genital System | 325.4 | 173.2 | 555.6 | 16 | 0.0 | 0.0 | 70.4 | 0 |
| Prostate | 325.4 | 173.2 | 555.6 | 16 | 0.0 | 0.0 | 70.4 | 0 |
| Testis | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Penis | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Urinary System | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Urinary Bladder | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Ureter | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | 31.9 | 15.4 | 57.5 | 11 |
| Retroperitoneum | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Respiratory System | 130.8 | 85.7 | 188.6 | 32 | 77.1 | 46.9 | 117.6 | 21 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Lung and Bronchus | 119.7 | 77.5 | 174.5 | 30 | 72.1 | 43.3 | 111.1 | 20 |
| Pleura | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Breast | 0.0 | 0.0 | 13.5 | 0 | 150.9 | 114.0 | 195.9 | 59 |
| Female Genital System | 0.0 | 0.0 | 13.5 | 0 | 22.6 | 9.5 | 44.8 | 8 |
| Cervix Uteri | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Ovary | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Vulva | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Male Genital System | 74.6 | 42.0 | 120.3 | 19 | 0.0 | 0.0 | 10.1 | 0 |
| Prostate | 67.3 | 35.9 | 112.2 | 16 | 0.0 | 0.0 | 10.1 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Urinary System | 44.4 | 21.4 | 79.8 | 13 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 24.3 | 10.2 | 49.1 | 8 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Larynx | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Bones and Joints | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| Breast | 0.0 | 0.0 | 45.6 | 0 | 89.3 | 48.0 | 155.2 | 14 |
| Female Genital System | 0.0 | 0.0 | 45.6 | 0 | 46.1 | 18.0 | 100.8 | 7 |
| Cervix Uteri | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Ovary | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Vulva | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Male Genital System | 97.0 | 31.9 | 211.4 | 6 | 0.0 | 0.0 | 35.2 | 0 |
| Prostate | 97.0 | 31.9 | 211.4 | 6 | 0.0 | 0.0 | 35.2 | 0 |
| Testis | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Penis | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 7.1 | 4.0 | 12.1 | 16 | 7.7 | 4.0 | 13.2 | 14 |
| Brain | 6.4 | 3.5 | 11.1 | 15 | 7.7 | 4.0 | 13.2 | 14 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Endocrine System | 7.2 | 4.0 | 12.4 | 16 | 8.4 | 5.1 | 13.2 | 20 |
| Thyroid | 6.4 | 3.4 | 11.4 | 14 | 8.4 | 5.1 | 13.2 | 20 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.3 | 0 |
| Lymphoma | 28.0 | 19.9 | 38.1 | 51 | 22.2 | 14.8 | 31.8 | 33 |
| Hodgkin Lymphoma | 3.9 | 1.4 | 8.6 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 3.9 | 1.4 | 8.6 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Non-Hodgkin Lymphoma | 24.1 | 16.6 | 33.7 | 44 | 20.1 | 13.0 | 29.4 | 29 |
| NHL - Nodal | 17.5 | 11.0 | 26.2 | 30 | 12.4 | 7.1 | 19.9 | 18 |
| NHL - Extranodal | 6.6 | 3.4 | 11.9 | 14 | 7.7 | 3.6 | 14.2 | 11 |
| Myeloma | 5.2 | 2.2 | 10.5 | 9 | 5.3 | 1.9 | 11.1 | 6 |
| Leukemia | 24.1 | 15.4 | 35.4 | 33 | 17.3 | 11.1 | 25.6 | 28 |
| Lymphocytic Leukemia | 9.2 | 4.8 | 15.9 | 16 | 7.5 | 3.7 | 13.3 | 12 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 8.1 | 3.9 | 14.6 | 13 | 4.2 | 1.5 | 9.1 | 6 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 14.8 | 7.8 | 24.8 | 17 | 8.9 | 4.5 | 15.5 | 14 |
| Acute Myeloid Leukemia | 10.2 | 4.6 | 18.9 | 12 | 6.8 | 2.9 | 12.9 | 10 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Leukemia | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 2.8 | 0 | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | 0.0 | 0.0 | 2.8 | 0 | 0.0 | 0.0 | 2.3 | 0 |
| Miscellaneous | 16.2 | 9.0 | 26.1 | 21 | 10.7 | 5.6 | 18.0 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Brain | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Endocrine System | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 70.4 | 0 |
| Myeloma | 0.0 | 0.0 | 110.2 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Mesothelioma | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 110.2 | 0 | 0.0 | 0.0 | 70.4 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 10.1 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Brain | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 13.5 | 0 | 14.7 | 5.8 | 31.6 | 7 |
| Thyroid | 0.0 | 0.0 | 13.5 | 0 | 14.7 | 5.8 | 31.6 | 7 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Lymphoma | 17.2 | 5.4 | 40.6 | 6 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Non-Hodgkin Lymphoma | 17.2 | 5.4 | 40.6 | 6 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 13.5 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Mesothelioma | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 13.5 | 0 | 0.0 | 0.0 | 10.1 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 45.6 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 45.6 | 0 | 0.0 | 0.0 | 35.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 35.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 619.9 | 574.8 | 667.4 | 944 | 446.5 | 413.2 | 481.6 | 764 |
| Oral Cavity and Pharynx | 16.8 | 10.8 | 24.8 | 32 | 4.6 | 1.9 | 9.2 | 8 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Tongue | 5.0 | 1.8 | 10.6 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Tonsil | 4.4 | 2.0 | 8.9 | 10 | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 2.7 | 0 | 0.0 | 0.0 | 2.2 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Digestive System | 104.5 | 86.3 | 125.2 | 155 | 81.8 | 66.9 | 98.7 | 117 |
| Esophagus | 7.4 | 3.5 | 13.7 | 13 | 5.5 | 2.1 | 11.2 | 7 |
| Stomach | 7.7 | 3.9 | 13.6 | 15 | 5.6 | 2.2 | 11.3 | 7 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Colon and Rectum | 70.5 | 55.1 | 88.4 | 93 | 43.2 | 32.6 | 56.0 | 61 |
| Colon excluding Rectum | 52.5 | 39.0 | 68.7 | 66 | 29.3 | 20.5 | 40.2 | 39 |
| Cecum | 7.5 | 3.6 | 13.8 | 11 | 10.6 | 5.7 | 17.6 | 15 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Ascending Colon | 15.8 | 8.0 | 26.8 | 13 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 3.7 | 1.1 | 8.7 | 6 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 14.0 | 7.9 | 22.6 | 20 | 5.8 | 2.4 | 11.3 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 18.0 | 11.2 | 27.0 | 27 | 14.0 | 8.5 | 21.5 | 22 |
| Rectosigmoid Junction | 4.8 | 1.6 | 10.6 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 13.2 | 7.7 | 21.0 | 21 | 13.1 | 7.8 | 20.6 | 20 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 8.2 | 3.8 | 15.0 | 15 | 6.0 | 2.6 | 11.5 | 10 |
| Liver | 6.6 | 2.6 | 13.4 | 11 | 5.3 | 2.1 | 10.7 | 8 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 696.8 | 461.4 | 1,004.3 | 37 | 480.6 | 286.5 | 741.0 | 27 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Lip | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Tongue | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Nasopharynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Tonsil | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Digestive System | 125.8 | 40.2 | 293.0 | 7 | 186.9 | 73.4 | 370.9 | 8 |
| Esophagus | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 161.4 | 55.2 | 342.2 | 6 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Transverse Colon | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 541.3 | 449.5 | 644.7 | 148 | 489.0 | 416.7 | 569.6 | 175 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 26.0 | 11.6 | 49.4 | 9 |
| Lip | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Tongue | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Oropharynx | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Hypopharynx | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Digestive System | 197.0 | 143.6 | 262.1 | 53 | 173.8 | 130.0 | 226.6 | 56 |
| Esophagus | 29.6 | 9.9 | 63.8 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 24.2 | 7.9 | 53.4 | 6 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 103.0 | 66.0 | 151.7 | 28 | 109.8 | 76.4 | 152.0 | 37 |
| Colon excluding Rectum | 48.3 | 25.0 | 83.1 | 13 | 90.2 | 59.7 | 129.6 | 29 |
| Cecum | ^ | ^ | ^ | ^ | 16.2 | 5.9 | 34.9 | 6 |
| Appendix | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 36.4 | 18.4 | 63.5 | 12 |
| Large Intestine, NOS | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 54.7 | 28.4 | 93.3 | 15 | 19.7 | 8.2 | 39.5 | 8 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 36.5 | 16.7 | 67.9 | 11 | 17.5 | 6.7 | 36.8 | 7 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 18.9 | 6.2 | 43.1 | 6 | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 371.1 | 243.4 | 536.9 | 32 | 272.5 | 181.0 | 391.5 | 35 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Tongue | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Nasopharynx | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Tonsil | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Oropharynx | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Hypopharynx | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Digestive System | 133.3 | 59.6 | 248.6 | 11 | 51.2 | 17.9 | 113.0 | 6 |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Small Intestine | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Appendix | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Ascending Colon | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Transverse Colon | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Gallbladder | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 5.7 | 2.6 | 11.0 | 11 | 13.8 | 8.1 | 21.5 | 19 |
| Retroperitoneum | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Respiratory System | 89.8 | 72.9 | 109.2 | 130 | 65.7 | 52.8 | 80.5 | 100 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 6.9 | 3.0 | 13.2 | 10 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 81.0 | 64.8 | 99.6 | 116 | 63.0 | 50.3 | 77.6 | 94 |
| Pleura | 0.0 | 0.0 | 2.7 | 0 | 0.0 | 0.0 | 2.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Bones and Joints | 2.5 | 1.0 | 5.9 | 7 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 17.6 | 11.8 | 25.2 | 37 | 18.7 | 12.6 | 26.8 | 34 |
| Melanoma of the Skin | 17.3 | 11.5 | 24.9 | 36 | 17.9 | 11.8 | 25.8 | 32 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 141.2 | 123.5 | 160.6 | 264 |
| Female Genital System | 0.0 | 0.0 | 2.7 | 0 | 45.9 | 36.0 | 57.4 | 85 |
| Cervix Uteri | 0.0 | 0.0 | 2.7 | 0 | 9.2 | 5.3 | 14.8 | 19 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.7 | 0 | 25.5 | 18.3 | 34.5 | 47 |
| Corpus Uteri | 0.0 | 0.0 | 2.7 | 0 | 24.5 | 17.5 | 33.2 | 46 |
| Uterus, NOS | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 2.7 | 0 | 8.1 | 4.3 | 13.9 | 14 |
| Vagina | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 2.7 | 0 | 0.0 | 0.0 | 2.2 | 0 |
| Male Genital System | 207.3 | 181.8 | 235.1 | 312 | 0.0 | 0.0 | 2.2 | 0 |
| Prostate | 199.4 | 174.2 | 226.9 | 292 | 0.0 | 0.0 | 2.2 | 0 |
| Testis | 7.4 | 4.4 | 12.2 | 19 | 0.0 | 0.0 | 2.2 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.7 | 0 | 0.0 | 0.0 | 2.2 | 0 |
| Urinary System | 77.3 | 61.5 | 95.5 | 109 | 16.0 | 10.1 | 23.8 | 26 |
| Urinary Bladder | 52.4 | 39.6 | 67.6 | 73 | 8.5 | 4.4 | 14.5 | 14 |
| Kidney and Renal Pelvis | 23.4 | 15.0 | 34.4 | 34 | 6.6 | 3.1 | 12.1 | 11 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Pancreas | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Larynx | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Breast | 0.0 | 0.0 | 101.1 | 0 | 151.3 | 53.2 | 318.9 | 9 |
| Female Genital System | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Ovary | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Vagina | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Vulva | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Male Genital System | 313.3 | 160.2 | 544.3 | 15 | 0.0 | 0.0 | 65.3 | 0 |
| Prostate | 313.3 | 160.2 | 544.3 | 15 | 0.0 | 0.0 | 65.3 | 0 |
| Testis | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Penis | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Ureter | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Respiratory System | 139.2 | 93.2 | 197.6 | 34 | 73.0 | 45.6 | 109.6 | 23 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Larynx | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 136.4 | 90.8 | 194.6 | 33 | 70.2 | 43.3 | 106.3 | 22 |
| Pleura | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Bones and Joints | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Breast | 0.0 | 0.0 | 13.1 | 0 | 126.6 | 92.9 | 168.3 | 50 |
| Female Genital System | 0.0 | 0.0 | 13.1 | 0 | 39.2 | 22.3 | 64.0 | 16 |
| Cervix Uteri | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 13.1 | 0 | 22.0 | 10.0 | 42.1 | 9 |
| Corpus Uteri | 0.0 | 0.0 | 13.1 | 0 | 22.0 | 10.0 | 42.1 | 9 |
| Uterus, NOS | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Ovary | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Vulva | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 101.7 | 63.4 | 152.5 | 28 | 0.0 | 0.0 | 9.6 | 0 |
| Prostate | 94.6 | 57.1 | 144.9 | 24 | 0.0 | 0.0 | 9.6 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Penis | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Urinary System | 43.7 | 23.0 | 75.0 | 13 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 23.8 | 9.4 | 49.2 | 7 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 19.9 | 7.2 | 43.6 | 6 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Larynx | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Breast | 0.0 | 0.0 | 42.5 | 0 | 79.0 | 39.6 | 142.8 | 12 |
| Female Genital System | 0.0 | 0.0 | 42.5 | 0 | 39.3 | 14.0 | 90.1 | 6 |
| Cervix Uteri | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Ovary | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Vagina | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Vulva | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Male Genital System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Prostate | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Testis | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Penis | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 42.5 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Brain and Other Nervous System | 13.5 | 7.6 | 21.8 | 23 | 5.2 | 2.4 | 9.9 | 10 |
| Brain | 13.5 | 7.6 | 21.8 | 23 | 4.0 | 1.6 | 8.2 | 8 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Endocrine System | 5.4 | 2.9 | 9.7 | 14 | 17.1 | 11.7 | 24.1 | 36 |
| Thyroid | 3.9 | 2.0 | 7.7 | 11 | 13.0 | 8.6 | 19.1 | 29 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 4.0 | 1.5 | 8.6 | 7 |
| Lymphoma | 29.0 | 20.3 | 39.9 | 50 | 20.9 | 14.2 | 29.6 | 35 |
| Hodgkin Lymphoma | 3.1 | 1.2 | 7.0 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 3.1 | 1.2 | 7.0 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.7 | 0 | 0.0 | 0.0 | 2.2 | 0 |
| Non-Hodgkin Lymphoma | 25.9 | 17.6 | 36.5 | 43 | 19.6 | 13.0 | 28.1 | 32 |
| NHL - Nodal | 16.7 | 10.7 | 24.8 | 31 | 14.9 | 9.3 | 22.5 | 25 |
| NHL - Extranodal | 9.2 | 4.1 | 17.1 | 12 | 4.7 | 1.8 | 9.7 | 7 |
| Myeloma | 6.2 | 2.6 | 12.1 | 9 | ^ | ^ | ^ | ^ |
| Leukemia | 15.0 | 8.4 | 24.2 | 21 | 11.4 | 6.7 | 18.0 | 20 |
| Lymphocytic Leukemia | 8.5 | 4.0 | 15.3 | 13 | 3.7 | 1.4 | 7.9 | 7 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 6.5 | 2.5 | 13.2 | 8 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Myeloid and Monocytic Leukemia | 5.4 | 1.6 | 12.4 | 6 | 6.6 | 3.2 | 11.9 | 12 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 3.7 | 1.2 | 8.3 | 6 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 2.7 | 0 | ^ | ^ | ^ | ^ |
| Miscellaneous | 25.2 | 16.1 | 37.1 | 33 | 12.1 | 6.9 | 19.5 | 18 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Brain | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |
| Myeloma | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 101.1 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 101.1 | 0 | 0.0 | 0.0 | 65.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 65.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Lymphoma | 22.6 | 9.6 | 45.7 | 8 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Non-Hodgkin Lymphoma | 19.8 | 7.8 | 42.0 | 7 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 13.1 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.6 | 0 |
| Mesothelioma | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 13.1 | 0 | 0.0 | 0.0 | 9.6 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Brain | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Mesothelioma | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 42.5 | 0 | 0.0 | 0.0 | 32.6 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 32.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 630.9 | 586.9 | 677.0 | 1,019 | 399.6 | 369.1 | 431.8 | 726 |
| Oral Cavity and Pharynx | 15.3 | 10.0 | 22.5 | 32 | 5.7 | 2.5 | 11.0 | 9 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Tongue | 3.9 | 1.6 | 8.2 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Digestive System | 103.4 | 85.9 | 123.1 | 165 | 67.0 | 54.0 | 81.9 | 104 |
| Esophagus | 10.5 | 5.3 | 18.3 | 15 | ^ | ^ | ^ | ^ |
| Stomach | 8.4 | 3.6 | 15.8 | 11 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 62.1 | 48.7 | 77.6 | 96 | 40.4 | 30.4 | 52.5 | 60 |
| Colon excluding Rectum | 40.2 | 29.2 | 53.6 | 57 | 30.8 | 22.1 | 41.5 | 45 |
| Cecum | 9.4 | 3.8 | 18.0 | 9 | 7.4 | 3.5 | 13.2 | 11 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Ascending Colon | 5.5 | 2.3 | 10.9 | 9 | 4.6 | 1.8 | 9.5 | 7 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 6.5 | 2.7 | 12.6 | 8 |
| Transverse Colon | 4.7 | 1.8 | 9.8 | 8 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Sigmoid Colon | 12.5 | 6.7 | 20.7 | 18 | 8.2 | 4.3 | 14.0 | 14 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 21.8 | 14.8 | 30.8 | 39 | 9.6 | 5.2 | 16.1 | 15 |
| Rectosigmoid Junction | 5.1 | 2.2 | 10.2 | 10 | ^ | ^ | ^ | ^ |
| Rectum | 16.7 | 10.6 | 24.8 | 29 | 7.9 | 3.8 | 14.1 | 11 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 2.9 | 1.0 | 6.7 | 6 |
| Liver and Intrahepatic Bile Duct | 5.5 | 2.6 | 10.4 | 12 | ^ | ^ | ^ | ^ |
| Liver | 5.1 | 2.3 | 9.9 | 11 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Gallbladder | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 512.0 | 339.8 | 742.0 | 36 | 398.5 | 240.9 | 611.6 | 26 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Tongue | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Tonsil | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Stomach | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Small Intestine | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Appendix | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Ascending Colon | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Rectum | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 510.7 | 425.0 | 607.2 | 154 | 500.3 | 425.3 | 583.9 | 172 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Tongue | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Digestive System | 203.7 | 145.7 | 274.4 | 49 | 171.1 | 126.8 | 224.7 | 53 |
| Esophagus | 27.9 | 9.4 | 60.0 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 53.0 | 24.7 | 95.3 | 12 | 32.0 | 14.7 | 58.7 | 10 |
| Small Intestine | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 59.6 | 32.5 | 98.3 | 16 | 107.5 | 73.1 | 151.2 | 33 |
| Colon excluding Rectum | 37.5 | 15.8 | 72.0 | 9 | 73.0 | 45.0 | 110.4 | 22 |
| Cecum | 0.0 | 0.0 | 12.5 | 0 | 23.7 | 8.7 | 49.2 | 6 |
| Appendix | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 22.1 | 8.8 | 45.8 | 7 | 34.5 | 16.7 | 61.6 | 11 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 34.5 | 16.7 | 61.6 | 11 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 375.1 | 239.7 | 551.1 | 32 | 343.1 | 244.0 | 467.0 | 50 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Lip | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Tongue | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Salivary Gland | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Tonsil | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Oropharynx | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | 73.8 | 29.4 | 147.4 | 10 |
| Esophagus | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Colon and Rectum | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Appendix | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Ascending Colon | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Transverse Colon | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Descending Colon | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Rectum | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Gallbladder | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 12.5 | 7.1 | 20.2 | 22 | 11.0 | 6.2 | 17.8 | 17 |
| Retroperitoneum | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 100.0 | 82.2 | 120.1 | 141 | 54.6 | 43.3 | 67.9 | 88 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 4.4 | 1.6 | 9.4 | 8 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 93.1 | 75.8 | 112.8 | 129 | 52.7 | 41.5 | 65.7 | 85 |
| Pleura | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 3.0 | 1.2 | 6.6 | 7 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 19.7 | 13.1 | 28.4 | 40 | 15.4 | 10.3 | 22.2 | 33 |
| Melanoma of the Skin | 17.4 | 11.2 | 25.6 | 36 | 15.0 | 10.0 | 21.7 | 32 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 131.0 | 114.5 | 149.3 | 255 |
| Female Genital System | 0.0 | 0.0 | 2.6 | 0 | 46.3 | 36.7 | 57.6 | 91 |
| Cervix Uteri | 0.0 | 0.0 | 2.6 | 0 | 8.5 | 4.9 | 13.7 | 19 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.6 | 0 | 18.2 | 12.2 | 26.0 | 33 |
| Corpus Uteri | 0.0 | 0.0 | 2.6 | 0 | 16.9 | 11.2 | 24.4 | 31 |
| Uterus, NOS | 0.0 | 0.0 | 2.6 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 2.6 | 0 | 13.3 | 8.5 | 20.0 | 26 |
| Vagina | 0.0 | 0.0 | 2.6 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.6 | 0 | 4.7 | 2.0 | 9.2 | 9 |
| Other Female Genital Organs | 0.0 | 0.0 | 2.6 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 221.6 | 196.2 | 249.2 | 355 | 0.0 | 0.0 | 2.1 | 0 |
| Prostate | 213.3 | 188.2 | 240.6 | 334 | 0.0 | 0.0 | 2.1 | 0 |
| Testis | 8.0 | 4.8 | 12.9 | 20 | 0.0 | 0.0 | 2.1 | 0 |
| Penis | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Urinary System | 78.2 | 62.8 | 95.8 | 118 | 14.4 | 8.9 | 21.9 | 24 |
| Urinary Bladder | 57.5 | 44.0 | 73.3 | 78 | 4.2 | 1.5 | 9.1 | 6 |
| Kidney and Renal Pelvis | 20.7 | 13.9 | 29.6 | 40 | 8.9 | 4.8 | 14.9 | 16 |
| Ureter | 0.0 | 0.0 | 2.6 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Larynx | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Breast | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 87.9 | 0 | 104.9 | 36.7 | 228.2 | 7 |
| Cervix Uteri | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Ovary | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Male Genital System | 217.9 | 121.4 | 370.2 | 16 | 0.0 | 0.0 | 61.3 | 0 |
| Prostate | 217.9 | 121.4 | 370.2 | 16 | 0.0 | 0.0 | 61.3 | 0 |
| Testis | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Penis | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Pancreas | 27.5 | 9.5 | 58.5 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 96.0 | 65.1 | 136.4 | 32 | 92.9 | 61.5 | 133.4 | 29 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 87.8 | 58.3 | 126.9 | 29 | 90.6 | 59.5 | 130.8 | 28 |
| Pleura | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 12.5 | 0 | 127.2 | 92.3 | 170.3 | 47 |
| Female Genital System | 0.0 | 0.0 | 12.5 | 0 | 25.8 | 11.9 | 48.0 | 10 |
| Cervix Uteri | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Ovary | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 99.5 | 64.2 | 145.8 | 31 | 0.0 | 0.0 | 9.3 | 0 |
| Prostate | 87.4 | 53.6 | 132.7 | 25 | 0.0 | 0.0 | 9.3 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Penis | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Urinary System | 38.5 | 19.8 | 67.4 | 13 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 31.1 | 14.9 | 57.1 | 11 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Pancreas | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | ^ | ^ | ^ | ^ | 66.6 | 23.2 | 141.4 | 8 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Larynx | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | 66.6 | 23.2 | 141.4 | 8 |
| Pleura | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Bones and Joints | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 40.6 | 0 | 87.3 | 47.8 | 148.2 | 15 |
| Female Genital System | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Ovary | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Vulva | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 220.3 | 110.8 | 377.7 | 14 | 0.0 | 0.0 | 29.1 | 0 |
| Prostate | 214.7 | 106.0 | 371.9 | 13 | 0.0 | 0.0 | 29.1 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Penis | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Brain and Other Nervous System | 7.8 | 4.6 | 12.7 | 19 | 5.6 | 2.7 | 10.2 | 11 |
| Brain | 7.4 | 4.3 | 12.3 | 18 | 5.6 | 2.7 | 10.2 | 11 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Endocrine System | 4.6 | 2.0 | 9.1 | 10 | 10.1 | 6.6 | 15.1 | 26 |
| Thyroid | 3.6 | 1.6 | 7.3 | 9 | 9.3 | 6.0 | 14.2 | 24 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 39.0 | 27.9 | 52.5 | 60 | 22.9 | 16.1 | 31.4 | 42 |
| Hodgkin Lymphoma | 2.2 | 0.8 | 5.4 | 6 | 4.0 | 1.9 | 7.7 | 10 |
| Hodgkin - Nodal | 2.2 | 0.8 | 5.4 | 6 | 4.0 | 1.9 | 7.7 | 10 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Non-Hodgkin Lymphoma | 36.8 | 25.9 | 50.2 | 54 | 18.8 | 12.6 | 27.0 | 32 |
| NHL - Nodal | 27.0 | 17.9 | 38.7 | 40 | 13.9 | 8.5 | 21.2 | 23 |
| NHL - Extranodal | 9.8 | 4.6 | 17.6 | 14 | 4.9 | 2.2 | 9.5 | 9 |
| Myeloma | 5.3 | 2.5 | 10.1 | 11 | 3.4 | 1.1 | 7.6 | 6 |
| Leukemia | 14.7 | 9.0 | 22.4 | 29 | 10.4 | 5.7 | 17.3 | 15 |
| Lymphocytic Leukemia | 7.9 | 3.6 | 14.6 | 13 | 7.8 | 3.8 | 14.0 | 11 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 6.3 | 2.4 | 12.9 | 9 | 6.9 | 3.1 | 12.9 | 9 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Myeloid and Monocytic Leukemia | 6.4 | 3.4 | 11.3 | 15 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 3.1 | 1.3 | 6.7 | 8 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 2.6 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | 3.4 | 1.2 | 7.7 | 7 | 0.0 | 0.0 | 2.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 2.6 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Miscellaneous | 14.2 | 8.0 | 22.9 | 23 | 8.7 | 4.6 | 14.8 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Eye and Orbit | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Brain | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 87.9 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 61.3 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 87.9 | 0 | 0.0 | 0.0 | 61.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 12.5 | 0 | 17.8 | 7.1 | 36.7 | 8 |
| Thyroid | 0.0 | 0.0 | 12.5 | 0 | 17.8 | 7.1 | 36.7 | 8 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 9.9 | 3.4 | 25.6 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 12.5 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 12.5 | 0 | 0.0 | 0.0 | 9.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 19.2 | 7.1 | 40.2 | 7 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Brain | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 40.6 | 0 | 34.7 | 11.8 | 81.1 | 6 |
| Thyroid | 0.0 | 0.0 | 40.6 | 0 | 34.7 | 11.8 | 81.1 | 6 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 40.6 | 0 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Mesothelioma | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 40.6 | 0 | 0.0 | 0.0 | 29.1 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 29.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 551.7 | 511.6 | 593.9 | 938 | 405.1 | 374.6 | 437.4 | 748 |
| Oral Cavity and Pharynx | 14.6 | 8.6 | 23.0 | 27 | 4.6 | 1.7 | 9.5 | 7 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Tonsil | 1.9 | 0.7 | 4.9 | 6 | 0.0 | 0.0 | 2.1 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Digestive System | 98.3 | 81.2 | 117.6 | 164 | 57.7 | 46.0 | 71.2 | 95 |
| Esophagus | 8.3 | 4.7 | 13.7 | 19 | ^ | ^ | ^ | ^ |
| Stomach | 5.8 | 2.6 | 10.9 | 11 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 53.9 | 41.3 | 68.8 | 84 | 33.2 | 24.6 | 43.6 | 56 |
| Colon excluding Rectum | 38.1 | 27.3 | 51.3 | 53 | 20.2 | 13.5 | 28.8 | 32 |
| Cecum | ^ | ^ | ^ | ^ | 6.7 | 3.0 | 12.6 | 9 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Ascending Colon | 7.5 | 3.1 | 14.4 | 10 | 4.2 | 1.5 | 9.0 | 6 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Descending Colon | 7.4 | 3.3 | 13.9 | 9 | 0.0 | 0.0 | 2.1 | 0 |
| Sigmoid Colon | 10.4 | 5.3 | 18.1 | 15 | 4.5 | 2.0 | 8.6 | 10 |
| Large Intestine, NOS | 5.0 | 1.5 | 11.1 | 7 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 15.8 | 9.9 | 23.8 | 31 | 13.0 | 8.1 | 19.8 | 24 |
| Rectosigmoid Junction | 7.2 | 3.1 | 13.7 | 11 | 3.5 | 1.2 | 7.7 | 6 |
| Rectum | 8.7 | 4.9 | 14.2 | 20 | 9.5 | 5.4 | 15.5 | 18 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 4.3 | 1.8 | 8.8 | 8 |
| Liver and Intrahepatic Bile Duct | 8.8 | 4.5 | 15.4 | 18 | ^ | ^ | ^ | ^ |
| Liver | 8.8 | 4.5 | 15.4 | 18 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 516.7 | 319.5 | 779.7 | 29 | 392.1 | 239.6 | 597.2 | 26 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Lip | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Tongue | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Salivary Gland | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Nasopharynx | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Tonsil | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Hypopharynx | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Digestive System | 203.0 | 84.8 | 393.2 | 10 | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Colon and Rectum | 141.6 | 46.9 | 310.3 | 7 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Appendix | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Transverse Colon | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 572.1 | 480.1 | 674.9 | 163 | 498.8 | 426.3 | 579.2 | 182 |
| Oral Cavity and Pharynx | 30.8 | 15.1 | 55.9 | 11 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Nasopharynx | 17.7 | 6.4 | 39.0 | 6 | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Digestive System | 197.9 | 145.5 | 261.3 | 56 | 131.6 | 95.7 | 175.8 | 47 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Colon and Rectum | 115.8 | 76.6 | 166.3 | 33 | 84.5 | 55.3 | 122.3 | 28 |
| Colon excluding Rectum | 89.6 | 54.1 | 137.3 | 23 | 79.2 | 51.0 | 116.2 | 26 |
| Cecum | ^ | ^ | ^ | ^ | 23.6 | 9.1 | 47.6 | 7 |
| Appendix | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 30.4 | 13.3 | 58.3 | 9 | 36.2 | 18.9 | 61.9 | 13 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Rectum and Rectosigmoid Junction | 26.2 | 12.4 | 49.1 | 10 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Rectum | 20.8 | 8.8 | 42.0 | 8 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 456.7 | 317.7 | 630.5 | 42 | 263.8 | 182.9 | 366.9 | 40 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Lip | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Tongue | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Salivary Gland | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Tonsil | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Oropharynx | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Hypopharynx | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Digestive System | 203.8 | 110.8 | 335.1 | 17 | 46.5 | 17.7 | 98.3 | 7 |
| Esophagus | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Appendix | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Transverse Colon | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Descending Colon | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Gallbladder | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 13.4 | 7.0 | 22.5 | 19 | 7.0 | 3.3 | 12.6 | 11 |
| Retroperitoneum | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 97.4 | 80.4 | 116.6 | 148 | 63.2 | 50.5 | 77.7 | 95 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Larynx | 6.5 | 3.1 | 11.9 | 12 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 90.0 | 73.5 | 108.7 | 133 | 62.3 | 49.7 | 76.9 | 93 |
| Pleura | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.9 | 2.0 | 10.1 | 9 | 2.9 | 1.0 | 6.5 | 6 |
| Skin excluding Basal and Squamous | 23.4 | 16.7 | 31.9 | 55 | 11.6 | 7.1 | 17.7 | 24 |
| Melanoma of the Skin | 20.8 | 14.6 | 28.7 | 50 | 9.6 | 5.7 | 15.1 | 21 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 136.3 | 119.4 | 154.8 | 272 |
| Female Genital System | 0.0 | 0.0 | 2.4 | 0 | 46.6 | 37.1 | 57.8 | 94 |
| Cervix Uteri | 0.0 | 0.0 | 2.4 | 0 | 4.6 | 2.1 | 8.7 | 10 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.4 | 0 | 23.1 | 16.5 | 31.3 | 46 |
| Corpus Uteri | 0.0 | 0.0 | 2.4 | 0 | 21.8 | 15.5 | 29.9 | 44 |
| Uterus, NOS | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 2.4 | 0 | 16.3 | 11.0 | 23.3 | 34 |
| Vagina | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Male Genital System | 162.9 | 142.1 | 185.9 | 278 | 0.0 | 0.0 | 2.1 | 0 |
| Prostate | 156.0 | 135.4 | 178.7 | 260 | 0.0 | 0.0 | 2.1 | 0 |
| Testis | 5.7 | 3.2 | 9.8 | 15 | 0.0 | 0.0 | 2.1 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Urinary System | 59.0 | 46.2 | 73.9 | 94 | 23.8 | 16.7 | 32.7 | 41 |
| Urinary Bladder | 44.1 | 32.9 | 57.6 | 63 | 12.4 | 7.4 | 19.3 | 21 |
| Kidney and Renal Pelvis | 14.9 | 9.4 | 22.4 | 31 | 10.7 | 6.2 | 17.0 | 19 |
| Ureter | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Larynx | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Bones and Joints | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Breast | ^ | ^ | ^ | ^ | 108.2 | 41.5 | 225.5 | 9 |
| Female Genital System | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Ovary | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Vulva | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Male Genital System | 174.5 | 69.1 | 349.9 | 9 | 0.0 | 0.0 | 58.2 | 0 |
| Prostate | 174.5 | 69.1 | 349.9 | 9 | 0.0 | 0.0 | 58.2 | 0 |
| Testis | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Penis | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 17.6 | 6.0 | 39.4 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 136.6 | 93.1 | 191.5 | 36 | 68.7 | 43.0 | 103.0 | 23 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Lung and Bronchus | 127.2 | 85.0 | 181.0 | 33 | 68.7 | 43.0 | 103.0 | 23 |
| Pleura | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Breast | 0.0 | 0.0 | 11.9 | 0 | 170.3 | 131.0 | 217.3 | 67 |
| Female Genital System | 0.0 | 0.0 | 11.9 | 0 | 35.9 | 17.7 | 63.3 | 12 |
| Cervix Uteri | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Ovary | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Vulva | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Male Genital System | 94.3 | 59.5 | 140.2 | 28 | 0.0 | 0.0 | 9.0 | 0 |
| Prostate | 84.8 | 51.2 | 130.1 | 23 | 0.0 | 0.0 | 9.0 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Penis | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Urinary System | 75.8 | 43.0 | 121.1 | 19 | 19.8 | 8.4 | 39.2 | 8 |
| Urinary Bladder | 35.6 | 13.1 | 72.3 | 7 | 0.0 | 0.0 | 9.0 | 0 |
| Kidney and Renal Pelvis | 37.3 | 17.2 | 68.5 | 11 | 19.8 | 8.4 | 39.2 | 8 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Bones and Joints | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Breast | 0.0 | 0.0 | 37.8 | 0 | 65.6 | 31.7 | 120.9 | 11 |
| Female Genital System | 0.0 | 0.0 | 37.8 | 0 | 45.9 | 18.8 | 94.5 | 8 |
| Cervix Uteri | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Ovary | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Vulva | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Male Genital System | 131.3 | 62.5 | 236.2 | 11 | 0.0 | 0.0 | 26.4 | 0 |
| Prostate | 131.3 | 62.5 | 236.2 | 11 | 0.0 | 0.0 | 26.4 | 0 |
| Testis | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Penis | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Ureter | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 7.0 | 3.7 | 12.2 | 16 | 6.7 | 3.2 | 11.9 | 12 |
| Brain | 6.2 | 3.1 | 11.2 | 14 | 6.2 | 2.9 | 11.4 | 11 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 6.2 | 3.3 | 10.9 | 15 | 14.3 | 9.9 | 20.2 | 35 |
| Thyroid | 4.8 | 2.5 | 8.8 | 13 | 14.3 | 9.9 | 20.2 | 35 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Lymphoma | 32.4 | 23.4 | 43.5 | 59 | 15.7 | 10.2 | 23.1 | 29 |
| Hodgkin Lymphoma | 6.0 | 3.0 | 10.8 | 14 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 6.0 | 3.0 | 10.8 | 14 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Non-Hodgkin Lymphoma | 26.4 | 18.1 | 36.9 | 45 | 14.0 | 8.7 | 21.1 | 25 |
| NHL - Nodal | 17.9 | 11.2 | 27.0 | 31 | 8.7 | 4.8 | 14.3 | 17 |
| NHL - Extranodal | 8.5 | 4.2 | 14.9 | 14 | 5.3 | 2.2 | 10.6 | 8 |
| Myeloma | 3.5 | 1.4 | 7.4 | 8 | 6.2 | 2.8 | 11.7 | 10 |
| Leukemia | 13.0 | 8.2 | 19.7 | 26 | 6.2 | 2.9 | 11.4 | 11 |
| Lymphocytic Leukemia | 6.7 | 3.5 | 11.7 | 14 | 3.4 | 1.2 | 7.6 | 6 |
| Acute Lymphocytic Leukemia | 2.3 | 0.8 | 5.6 | 6 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Myeloid and Monocytic Leukemia | 5.9 | 2.8 | 11.1 | 11 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 2.6 | 0.9 | 6.1 | 6 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 2.4 | 0 | ^ | ^ | ^ | ^ |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 2.4 | 0 | 0.0 | 0.0 | 2.1 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | 6.2 | 2.1 | 13.4 | 6 | 0.0 | 0.0 | 2.1 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.1 | 0 |
| Miscellaneous | 19.5 | 11.8 | 29.6 | 26 | 8.7 | 4.6 | 14.7 | 15 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Myeloma | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Leukemia | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 77.3 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 77.3 | 0 | 0.0 | 0.0 | 58.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 58.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Myeloma | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 9.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 11.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 11.9 | 0 | 0.0 | 0.0 | 9.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Endocrine System | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Lymphoma | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 37.8 | 0 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Mesothelioma | 0.0 | 0.0 | 37.8 | 0 | 0.0 | 0.0 | 26.4 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 26.4 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 564.5 | 525.4 | 605.6 | 1,001 | 401.7 | 371.8 | 433.2 | 768 |
| Oral Cavity and Pharynx | 18.8 | 12.5 | 27.1 | 37 | 4.3 | 1.7 | 8.7 | 8 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Tongue | 3.5 | 1.5 | 7.2 | 9 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 4.5 | 1.4 | 10.0 | 7 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Tonsil | 1.8 | 0.7 | 4.7 | 6 | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Digestive System | 108.6 | 91.3 | 127.9 | 182 | 62.6 | 50.6 | 76.5 | 107 |
| Esophagus | 8.4 | 4.6 | 14.0 | 16 | ^ | ^ | ^ | ^ |
| Stomach | 5.9 | 2.6 | 11.3 | 9 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 59.2 | 46.3 | 74.3 | 95 | 32.5 | 24.0 | 42.8 | 56 |
| Colon excluding Rectum | 43.7 | 32.3 | 57.4 | 63 | 26.7 | 19.1 | 36.2 | 45 |
| Cecum | 10.4 | 5.0 | 18.3 | 14 | 7.7 | 3.7 | 13.8 | 11 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 5.0 | 1.9 | 10.1 | 8 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 4.4 | 1.5 | 9.7 | 6 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 15.8 | 9.1 | 25.0 | 22 | 5.3 | 2.5 | 9.8 | 11 |
| Large Intestine, NOS | 3.0 | 1.0 | 6.9 | 6 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 15.5 | 10.1 | 22.7 | 32 | 5.8 | 2.7 | 10.8 | 11 |
| Rectosigmoid Junction | 3.6 | 1.2 | 8.2 | 7 | ^ | ^ | ^ | ^ |
| Rectum | 11.9 | 7.4 | 18.2 | 25 | 5.4 | 2.4 | 10.4 | 10 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 7.4 | 4.1 | 12.4 | 18 | 5.1 | 2.0 | 10.1 | 8 |
| Liver | 7.0 | 3.8 | 11.9 | 17 | 4.2 | 1.6 | 8.9 | 7 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 805.1 | 534.8 | 1,145.9 | 39 | 318.8 | 201.3 | 478.3 | 27 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Lip | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Tongue | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Nasopharynx | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Tonsil | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Hypopharynx | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Digestive System | 198.1 | 76.2 | 395.4 | 9 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Small Intestine | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Ascending Colon | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Transverse Colon | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Descending Colon | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 560.6 | 474.7 | 656.4 | 174 | 600.0 | 522.1 | 685.6 | 227 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 16.5 | 5.7 | 35.9 | 6 |
| Lip | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Tongue | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Hypopharynx | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Digestive System | 218.8 | 165.6 | 282.3 | 67 | 185.1 | 141.7 | 236.7 | 65 |
| Esophagus | ^ | ^ | ^ | ^ | 19.6 | 7.1 | 41.2 | 6 |
| Stomach | 45.4 | 22.6 | 79.2 | 14 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 88.1 | 56.9 | 129.1 | 28 | 123.2 | 88.9 | 165.7 | 45 |
| Colon excluding Rectum | 74.5 | 45.7 | 113.5 | 23 | 101.1 | 70.4 | 139.8 | 37 |
| Cecum | ^ | ^ | ^ | ^ | 28.7 | 13.5 | 52.4 | 10 |
| Appendix | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 35.3 | 16.8 | 64.1 | 11 | 26.3 | 12.8 | 47.4 | 11 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 22.2 | 9.0 | 43.9 | 8 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 16.1 | 5.5 | 35.2 | 6 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Liver and Intrahepatic Bile Duct | 31.4 | 11.9 | 63.5 | 8 | ^ | ^ | ^ | ^ |
| Liver | 28.9 | 10.2 | 60.8 | 7 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 346.0 | 229.5 | 495.5 | 36 | 240.3 | 168.3 | 332.2 | 39 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Lip | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Tongue | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Tonsil | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Digestive System | 69.3 | 31.6 | 134.1 | 10 | 75.3 | 36.5 | 135.6 | 11 |
| Esophagus | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Stomach | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 55.0 | 21.1 | 117.6 | 7 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Appendix | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Transverse Colon | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Gallbladder | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 14.6 | 8.9 | 22.5 | 25 | 10.2 | 5.7 | 16.8 | 16 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 5.1 | 1.9 | 10.6 | 7 | ^ | ^ | ^ | ^ |
| Respiratory System | 89.6 | 73.6 | 107.8 | 141 | 59.5 | 47.7 | 73.2 | 98 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Larynx | 5.1 | 2.3 | 10.0 | 10 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 84.1 | 68.5 | 101.9 | 130 | 58.2 | 46.5 | 71.7 | 96 |
| Pleura | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.4 | 1.9 | 8.8 | 10 | 2.9 | 1.0 | 6.5 | 6 |
| Skin excluding Basal and Squamous | 16.9 | 11.2 | 24.4 | 35 | 14.6 | 9.6 | 21.3 | 30 |
| Melanoma of the Skin | 14.4 | 9.2 | 21.4 | 30 | 14.3 | 9.3 | 20.9 | 29 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 2.3 | 0 | 118.0 | 102.7 | 134.9 | 246 |
| Female Genital System | 0.0 | 0.0 | 2.3 | 0 | 49.9 | 40.0 | 61.4 | 100 |
| Cervix Uteri | 0.0 | 0.0 | 2.3 | 0 | 6.7 | 3.6 | 11.4 | 15 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.3 | 0 | 24.3 | 17.6 | 32.6 | 50 |
| Corpus Uteri | 0.0 | 0.0 | 2.3 | 0 | 24.0 | 17.3 | 32.2 | 49 |
| Uterus, NOS | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 2.3 | 0 | 10.4 | 6.3 | 16.1 | 22 |
| Vagina | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.3 | 0 | 3.9 | 1.4 | 8.4 | 6 |
| Other Female Genital Organs | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 172.0 | 151.4 | 194.4 | 320 | 0.0 | 0.0 | 2.0 | 0 |
| Prostate | 164.7 | 144.5 | 186.9 | 302 | 0.0 | 0.0 | 2.0 | 0 |
| Testis | 5.8 | 3.2 | 9.9 | 15 | 0.0 | 0.0 | 2.0 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Urinary System | 71.2 | 57.5 | 87.0 | 119 | 19.3 | 13.1 | 27.3 | 34 |
| Urinary Bladder | 47.9 | 36.8 | 61.1 | 79 | 6.1 | 2.9 | 11.2 | 11 |
| Kidney and Renal Pelvis | 23.0 | 15.6 | 32.5 | 39 | 11.2 | 6.6 | 17.5 | 20 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Pancreas | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Respiratory System | 165.4 | 57.1 | 349.0 | 7 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Larynx | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Lung and Bronchus | 165.4 | 57.1 | 349.0 | 7 | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Bones and Joints | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 68.7 | 0 | 150.7 | 76.9 | 266.4 | 13 |
| Female Genital System | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Ovary | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Vulva | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Male Genital System | 297.4 | 148.3 | 515.3 | 15 | 0.0 | 0.0 | 54.2 | 0 |
| Prostate | 297.4 | 148.3 | 515.3 | 15 | 0.0 | 0.0 | 54.2 | 0 |
| Testis | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Penis | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Urinary Bladder | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Ureter | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 25.6 | 9.9 | 52.1 | 7 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Respiratory System | 141.0 | 99.5 | 192.7 | 43 | 102.4 | 70.3 | 142.9 | 34 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Lung and Bronchus | 127.9 | 88.6 | 177.3 | 39 | 102.4 | 70.3 | 142.9 | 34 |
| Pleura | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Bones and Joints | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Breast | 0.0 | 0.0 | 11.5 | 0 | 156.7 | 119.8 | 201.0 | 64 |
| Female Genital System | 0.0 | 0.0 | 11.5 | 0 | 36.9 | 20.9 | 60.0 | 16 |
| Cervix Uteri | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 11.5 | 0 | 16.4 | 6.5 | 33.9 | 7 |
| Corpus Uteri | 0.0 | 0.0 | 11.5 | 0 | 16.4 | 6.5 | 33.9 | 7 |
| Uterus, NOS | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Ovary | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Male Genital System | 79.4 | 48.7 | 120.7 | 23 | 0.0 | 0.0 | 8.7 | 0 |
| Prostate | 70.9 | 42.1 | 110.2 | 20 | 0.0 | 0.0 | 8.7 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Urinary System | 56.9 | 32.4 | 91.6 | 17 | 29.5 | 14.4 | 52.9 | 11 |
| Urinary Bladder | 27.3 | 11.3 | 53.9 | 8 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 26.0 | 10.8 | 51.4 | 8 | 16.9 | 6.7 | 34.8 | 7 |
| Ureter | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Larynx | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Breast | 0.0 | 0.0 | 35.4 | 0 | 42.5 | 18.1 | 86.3 | 8 |
| Female Genital System | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Ovary | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Male Genital System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Prostate | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Testis | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Penis | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 8.7 | 5.0 | 14.2 | 20 | 5.0 | 2.1 | 9.9 | 8 |
| Brain | 8.3 | 4.7 | 13.7 | 19 | 4.1 | 1.4 | 8.8 | 6 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 4.0 | 2.0 | 7.6 | 11 | 21.0 | 15.7 | 27.7 | 54 |
| Thyroid | 4.0 | 2.0 | 7.6 | 11 | 20.3 | 15.1 | 26.8 | 52 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 2.3 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | 26.6 | 18.9 | 36.3 | 52 | 17.3 | 11.4 | 25.1 | 30 |
| Hodgkin Lymphoma | 4.1 | 1.9 | 7.8 | 10 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 4.1 | 1.9 | 7.8 | 10 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Non-Hodgkin Lymphoma | 22.5 | 15.3 | 31.9 | 42 | 15.6 | 9.9 | 23.2 | 26 |
| NHL - Nodal | 18.6 | 11.9 | 27.4 | 34 | 10.8 | 6.3 | 17.3 | 19 |
| NHL - Extranodal | 3.9 | 1.6 | 8.1 | 8 | 4.8 | 1.8 | 9.7 | 7 |
| Myeloma | 9.1 | 4.7 | 15.7 | 15 | 5.6 | 2.5 | 10.6 | 10 |
| Leukemia | 16.9 | 10.3 | 25.8 | 28 | 6.4 | 3.0 | 11.6 | 11 |
| Lymphocytic Leukemia | 12.4 | 6.7 | 20.5 | 19 | 5.0 | 2.2 | 9.7 | 9 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 8.6 | 3.8 | 16.0 | 12 | 4.1 | 1.5 | 8.6 | 7 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Myeloid and Monocytic Leukemia | 4.1 | 1.5 | 8.8 | 8 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.0 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | 0.0 | 0.0 | 2.3 | 0 | 0.0 | 0.0 | 2.0 | 0 |
| Miscellaneous | 13.2 | 8.0 | 20.3 | 24 | 10.0 | 5.6 | 16.3 | 17 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Brain | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 68.7 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 68.7 | 0 | 0.0 | 0.0 | 54.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 54.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Eye and Orbit | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 19.1 | 6.5 | 42.3 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 11.5 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Other Leukemia | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 11.5 | 0 | 0.0 | 0.0 | 8.7 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.7 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 15.0 | 5.3 | 33.0 | 6 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2004

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 24.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 35.4 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 35.4 | 0 | 0.0 | 0.0 | 24.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 485.1 | 449.9 | 522.1 | 909 | 442.7 | 411.3 | 475.8 | 838 |
| Oral Cavity and Pharynx | 12.7 | 8.4 | 18.5 | 31 | 4.8 | 2.2 | 9.2 | 10 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 3.8 | 1.6 | 7.8 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Digestive System | 100.3 | 83.9 | 118.7 | 174 | 67.0 | 54.4 | 81.3 | 110 |
| Esophagus | 10.5 | 6.1 | 16.7 | 21 | ^ | ^ | ^ | ^ |
| Stomach | 7.6 | 4.0 | 13.0 | 16 | ^ | ^ | ^ | ^ |
| Small Intestine | 3.9 | 1.6 | 8.1 | 8 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 49.5 | 38.3 | 62.7 | 86 | 40.3 | 30.6 | 51.8 | 64 |
| Colon excluding Rectum | 40.0 | 29.9 | 52.0 | 70 | 31.7 | 23.0 | 42.2 | 48 |
| Cecum | 13.1 | 7.2 | 21.3 | 18 | 7.6 | 3.8 | 13.2 | 12 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 5.7 | 2.6 | 10.6 | 11 | 9.0 | 4.6 | 15.5 | 12 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Descending Colon | 3.8 | 1.5 | 8.0 | 8 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 8.1 | 4.5 | 13.4 | 18 | 5.5 | 2.3 | 10.6 | 9 |
| Large Intestine, NOS | 5.2 | 1.8 | 11.0 | 8 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 9.5 | 5.1 | 16.1 | 16 | 8.6 | 4.7 | 14.3 | 16 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 7.8 | 3.8 | 13.9 | 13 | 6.7 | 3.3 | 11.9 | 12 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 10.2 | 5.2 | 17.5 | 15 | 0.0 | 0.0 | 1.9 | 0 |
| Liver | 9.4 | 4.6 | 16.6 | 14 | 0.0 | 0.0 | 1.9 | 0 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Gallbladder | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 511.3 | 339.8 | 733.5 | 39 | 346.0 | 208.0 | 532.5 | 23 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Lip | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Tongue | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Salivary Gland | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Nasopharynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Tonsil | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Hypopharynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Digestive System | 67.7 | 28.7 | 150.1 | 8 | 112.5 | 38.5 | 240.7 | 6 |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Small Intestine | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Ascending Colon | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Transverse Colon | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Descending Colon | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 537.2 | 454.9 | 629.1 | 181 | 502.3 | 432.4 | 579.8 | 197 |
| Oral Cavity and Pharynx | 24.4 | 11.5 | 45.9 | 10 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Oropharynx | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Digestive System | 210.3 | 159.2 | 271.3 | 67 | 123.7 | 89.6 | 165.6 | 46 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 36.8 | 17.1 | 67.1 | 12 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Colon and Rectum | 116.8 | 80.2 | 163.3 | 37 | 86.1 | 57.8 | 122.4 | 31 |
| Colon excluding Rectum | 76.0 | 46.4 | 116.0 | 23 | 70.5 | 44.9 | 104.2 | 25 |
| Cecum | ^ | ^ | ^ | ^ | 18.2 | 7.2 | 36.9 | 7 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 17.7 | 6.2 | 37.9 | 6 |
| Hepatic Flexure | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 20.2 | 7.8 | 41.1 | 7 |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 21.8 | 7.1 | 48.0 | 7 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Rectum and Rectosigmoid Junction | 40.8 | 21.6 | 69.3 | 14 | 15.6 | 5.6 | 33.6 | 6 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 27.6 | 12.7 | 51.9 | 10 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 381.9 | 257.4 | 539.3 | 35 | 321.0 | 241.7 | 418.0 | 60 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Lip | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Tongue | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Salivary Gland | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Nasopharynx | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Tonsil | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Oropharynx | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Hypopharynx | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Digestive System | 156.3 | 83.8 | 260.4 | 15 | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Small Intestine | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Colon and Rectum | 79.1 | 29.8 | 161.9 | 7 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Appendix | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Ascending Colon | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Descending Colon | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Gallbladder | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 9.3 | 4.6 | 16.2 | 15 | 13.2 | 8.0 | 20.2 | 21 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 3.5 | 1.2 | 7.7 | 6 | 3.3 | 1.1 | 7.3 | 6 |
| Respiratory System | 78.5 | 64.4 | 94.5 | 137 | 76.0 | 62.6 | 91.4 | 122 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 4.3 | 1.8 | 8.7 | 9 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 73.8 | 60.1 | 89.5 | 127 | 73.7 | 60.4 | 88.9 | 117 |
| Pleura | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.1 | 1.7 | 8.4 | 9 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 13.8 | 8.6 | 20.9 | 29 | 10.3 | 6.5 | 15.7 | 24 |
| Melanoma of the Skin | 9.8 | 6.0 | 15.3 | 24 | 9.1 | 5.5 | 14.3 | 21 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 136.4 | 119.7 | 154.6 | 278 |
| Female Genital System | 0.0 | 0.0 | 2.2 | 0 | 41.0 | 32.5 | 51.1 | 90 |
| Cervix Uteri | 0.0 | 0.0 | 2.2 | 0 | 8.7 | 5.2 | 13.8 | 20 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.2 | 0 | 16.7 | 11.4 | 23.6 | 37 |
| Corpus Uteri | 0.0 | 0.0 | 2.2 | 0 | 16.7 | 11.4 | 23.6 | 37 |
| Uterus, NOS | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Ovary | 0.0 | 0.0 | 2.2 | 0 | 12.3 | 7.8 | 18.4 | 26 |
| Vagina | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 123.4 | 106.7 | 141.9 | 245 | 0.0 | 0.0 | 1.9 | 0 |
| Prostate | 116.3 | 99.9 | 134.5 | 225 | 0.0 | 0.0 | 1.9 | 0 |
| Testis | 6.6 | 3.9 | 10.7 | 18 | 0.0 | 0.0 | 1.9 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Urinary System | 63.5 | 51.4 | 77.4 | 116 | 24.9 | 17.6 | 33.9 | 43 |
| Urinary Bladder | 41.5 | 31.6 | 53.3 | 71 | 12.1 | 7.2 | 18.8 | 20 |
| Kidney and Renal Pelvis | 21.0 | 14.7 | 29.1 | 44 | 11.3 | 6.6 | 17.8 | 20 |
| Ureter | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Pancreas | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Larynx | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Bones and Joints | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Breast | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Ovary | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Vulva | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Male Genital System | 295.2 | 161.4 | 485.9 | 20 | 0.0 | 0.0 | 51.2 | 0 |
| Prostate | 290.9 | 157.6 | 481.5 | 19 | 0.0 | 0.0 | 51.2 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Penis | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Ureter | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 24.7 | 8.1 | 53.7 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Respiratory System | 99.7 | 64.6 | 145.3 | 30 | 104.5 | 72.5 | 144.6 | 36 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 99.7 | 64.6 | 145.3 | 30 | 99.5 | 68.3 | 139.0 | 34 |
| Pleura | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 10.9 | 0 | 136.4 | 103.1 | 176.6 | 59 |
| Female Genital System | 0.0 | 0.0 | 10.9 | 0 | 47.4 | 27.8 | 74.7 | 19 |
| Cervix Uteri | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Ovary | 0.0 | 0.0 | 10.9 | 0 | 18.4 | 7.0 | 38.0 | 7 |
| Vagina | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Vulva | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 58.6 | 36.1 | 89.7 | 23 | 0.0 | 0.0 | 8.5 | 0 |
| Prostate | 54.3 | 32.4 | 85.0 | 20 | 0.0 | 0.0 | 8.5 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Penis | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Urinary System | 51.1 | 29.2 | 82.3 | 18 | 17.8 | 6.2 | 38.1 | 6 |
| Urinary Bladder | 27.2 | 11.2 | 53.7 | 8 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 23.9 | 11.1 | 45.3 | 10 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 96.6 | 36.8 | 194.2 | 7 | 50.1 | 17.8 | 106.2 | 6 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Larynx | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Lung and Bronchus | 96.6 | 36.8 | 194.2 | 7 | 50.1 | 17.8 | 106.2 | 6 |
| Pleura | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Bones and Joints | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Breast | 0.0 | 0.0 | 32.3 | 0 | 78.0 | 42.8 | 131.3 | 15 |
| Female Genital System | 0.0 | 0.0 | 32.3 | 0 | 80.8 | 46.4 | 132.2 | 17 |
| Cervix Uteri | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 32.3 | 0 | 46.4 | 22.2 | 87.9 | 10 |
| Corpus Uteri | 0.0 | 0.0 | 32.3 | 0 | 46.4 | 22.2 | 87.9 | 10 |
| Uterus, NOS | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Ovary | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Vulva | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Male Genital System | 66.6 | 24.8 | 139.2 | 7 | 0.0 | 0.0 | 22.5 | 0 |
| Prostate | 66.6 | 24.8 | 139.2 | 7 | 0.0 | 0.0 | 22.5 | 0 |
| Testis | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Penis | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.5 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Brain and Other Nervous System | 8.7 | 4.2 | 15.5 | 15 | 7.7 | 4.1 | 12.9 | 16 |
| Brain | 8.5 | 4.0 | 15.2 | 14 | 7.7 | 4.1 | 12.9 | 16 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Endocrine System | 6.5 | 3.8 | 10.7 | 18 | 15.9 | 10.8 | 22.4 | 35 |
| Thyroid | 6.2 | 3.6 | 10.3 | 17 | 15.9 | 10.8 | 22.4 | 35 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Lymphoma | 25.0 | 17.7 | 34.3 | 51 | 28.1 | 20.5 | 37.3 | 52 |
| Hodgkin Lymphoma | 3.3 | 1.5 | 6.7 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 3.3 | 1.5 | 6.7 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Non-Hodgkin Lymphoma | 21.7 | 14.7 | 30.7 | 42 | 26.4 | 19.1 | 35.4 | 49 |
| NHL - Nodal | 15.6 | 9.7 | 23.6 | 29 | 20.6 | 14.3 | 28.7 | 38 |
| NHL - Extranodal | 6.1 | 2.9 | 11.2 | 13 | 5.7 | 2.7 | 10.6 | 11 |
| Myeloma | 7.4 | 3.7 | 13.2 | 12 | 3.0 | 1.0 | 6.7 | 6 |
| Leukemia | 18.8 | 12.3 | 27.3 | 34 | 12.2 | 7.8 | 18.3 | 26 |
| Lymphocytic Leukemia | 13.9 | 8.2 | 21.6 | 24 | 6.4 | 3.3 | 11.2 | 13 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 10.9 | 5.9 | 18.1 | 18 | 4.6 | 2.0 | 9.0 | 9 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Myeloid and Monocytic Leukemia | 5.0 | 2.1 | 9.7 | 10 | 5.5 | 2.7 | 10.0 | 12 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 4.4 | 1.9 | 8.7 | 9 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Other Leukemia | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 2.2 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 2.2 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Mesothelioma | 4.2 | 1.3 | 9.6 | 6 | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Miscellaneous | 15.8 | 9.3 | 24.6 | 25 | 10.3 | 5.7 | 16.8 | 16 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Brain | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 64.2 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 64.2 | 0 | 0.0 | 0.0 | 51.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 51.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Endocrine System | ^ | ^ | ^ | ^ | 12.0 | 4.2 | 26.9 | 6 |
| Thyroid | ^ | ^ | ^ | ^ | 12.0 | 4.2 | 26.9 | 6 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Lymphoma | 24.3 | 8.1 | 52.7 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Non-Hodgkin Lymphoma | 24.3 | 8.1 | 52.7 | 7 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 16.0 | 5.2 | 36.5 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.5 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Other Leukemia | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 10.9 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Mesothelioma | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 10.9 | 0 | 0.0 | 0.0 | 8.5 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2005

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Brain | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Endocrine System | 0.0 | 0.0 | 32.3 | 0 | 30.8 | 10.7 | 70.0 | 6 |
| Thyroid | 0.0 | 0.0 | 32.3 | 0 | 30.8 | 10.7 | 70.0 | 6 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Other Leukemia | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 32.3 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Mesothelioma | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 32.3 | 0 | 0.0 | 0.0 | 22.5 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 501.2 | 466.0 | 538.3 | 976 | 414.9 | 385.4 | 445.9 | 843 |
| Oral Cavity and Pharynx | 15.7 | 10.4 | 22.8 | 34 | 5.5 | 2.4 | 10.4 | 9 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 4.0 | 1.5 | 8.3 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Tonsil | 5.8 | 3.0 | 10.1 | 15 | 0.0 | 0.0 | 1.9 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Digestive System | 92.8 | 77.5 | 110.1 | 173 | 73.1 | 60.5 | 87.5 | 135 |
| Esophagus | 8.9 | 4.7 | 15.0 | 18 | 3.3 | 1.1 | 7.4 | 6 |
| Stomach | 5.9 | 2.3 | 11.8 | 10 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 47.6 | 37.0 | 60.1 | 89 | 37.6 | 28.8 | 48.0 | 71 |
| Colon excluding Rectum | 35.6 | 26.5 | 46.5 | 63 | 29.9 | 22.1 | 39.5 | 55 |
| Cecum | 11.8 | 6.6 | 19.0 | 18 | 9.4 | 5.3 | 15.2 | 18 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 4.2 | 1.5 | 8.9 | 7 | 8.8 | 4.6 | 14.8 | 14 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 10.3 | 6.0 | 16.5 | 20 | 5.7 | 2.7 | 10.5 | 11 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 3.4 | 1.2 | 7.4 | 6 |
| Rectum and Rectosigmoid Junction | 12.1 | 7.1 | 18.9 | 26 | 7.6 | 4.2 | 12.7 | 16 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 10.2 | 5.7 | 16.7 | 22 | 5.3 | 2.6 | 9.5 | 12 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 4.4 | 2.0 | 8.5 | 10 |
| Liver and Intrahepatic Bile Duct | 9.4 | 4.8 | 16.3 | 16 | 3.6 | 1.4 | 7.4 | 8 |
| Liver | 9.4 | 4.8 | 16.3 | 16 | 3.3 | 1.2 | 7.1 | 7 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 442.8 | 279.9 | 658.9 | 32 | 214.3 | 114.8 | 358.5 | 16 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Lip | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Tongue | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Salivary Gland | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Nasopharynx | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Tonsil | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Oropharynx | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hypopharynx | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Digestive System | 170.9 | 70.8 | 332.5 | 11 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Stomach | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Appendix | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Ascending Colon | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Transverse Colon | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Gallbladder | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Alaska Native/American Indian | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 406.7 | 339.8 | 482.3 | 147 | 471.0 | 405.7 | 543.2 | 197 |
| Oral Cavity and Pharynx | 14.3 | 5.2 | 32.0 | 6 | 16.8 | 7.1 | 33.5 | 8 |
| Lip | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 10.6 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 10.6 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Digestive System | 149.9 | 109.4 | 199.5 | 52 | 133.8 | 99.1 | 176.0 | 52 |
| Esophagus | 19.7 | 7.5 | 41.4 | 7 | ^ | ^ | ^ | ^ |
| Stomach | 12.7 | 4.6 | 29.0 | 6 | 16.7 | 6.5 | 34.5 | 7 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Colon and Rectum | 92.0 | 60.0 | 133.8 | 30 | 75.3 | 50.4 | 107.7 | 30 |
| Colon excluding Rectum | 70.6 | 42.4 | 108.8 | 22 | 64.8 | 41.6 | 95.7 | 25 |
| Cecum | ^ | ^ | ^ | ^ | 19.3 | 7.7 | 38.9 | 7 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 25.2 | 11.0 | 48.7 | 9 | 16.6 | 6.6 | 34.0 | 7 |
| Large Intestine, NOS | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Rectum and Rectosigmoid Junction | 21.5 | 8.7 | 43.5 | 8 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Rectum | 17.0 | 5.8 | 38.0 | 6 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 10.6 | 0 | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 10.6 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 403.2 | 283.7 | 551.5 | 44 | 337.2 | 254.2 | 438.1 | 60 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Tongue | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Salivary Gland | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Nasopharynx | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Hypopharynx | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Digestive System | 120.2 | 63.9 | 203.4 | 15 | 59.4 | 27.1 | 111.5 | 10 |
| Esophagus | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Small Intestine | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 52.7 | 17.1 | 118.0 | 6 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Appendix | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Transverse Colon | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 12.7 | 7.6 | 19.8 | 24 | 11.0 | 6.3 | 17.6 | 18 |
| Retroperitoneum | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 91.5 | 76.6 | 108.2 | 170 | 62.3 | 50.6 | 75.7 | 109 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Larynx | 7.8 | 4.1 | 13.3 | 15 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 81.7 | 67.7 | 97.6 | 152 | 61.3 | 49.7 | 74.7 | 106 |
| Pleura | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.3 | 2.0 | 8.4 | 10 | 2.1 | 0.8 | 4.9 | 6 |
| Skin excluding Basal and Squamous | 13.9 | 8.6 | 21.1 | 29 | 13.3 | 8.8 | 19.4 | 30 |
| Melanoma of the Skin | 11.6 | 7.2 | 17.6 | 27 | 13.3 | 8.8 | 19.4 | 30 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Breast | 0.0 | 0.0 | 2.1 | 0 | 129.2 | 113.4 | 146.6 | 277 |
| Female Genital System | 0.0 | 0.0 | 2.1 | 0 | 43.4 | 34.9 | 53.4 | 101 |
| Cervix Uteri | 0.0 | 0.0 | 2.1 | 0 | 6.3 | 3.5 | 10.5 | 16 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 2.1 | 0 | 24.2 | 17.9 | 32.1 | 55 |
| Corpus Uteri | 0.0 | 0.0 | 2.1 | 0 | 24.2 | 17.9 | 32.1 | 55 |
| Uterus, NOS | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Ovary | 0.0 | 0.0 | 2.1 | 0 | 9.3 | 5.8 | 14.4 | 23 |
| Vagina | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 137.9 | 120.5 | 157.0 | 290 | 0.0 | 0.0 | 1.9 | 0 |
| Prostate | 131.1 | 114.0 | 150.0 | 271 | 0.0 | 0.0 | 1.9 | 0 |
| Testis | 5.7 | 3.3 | 9.5 | 17 | 0.0 | 0.0 | 1.9 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Urinary System | 65.0 | 52.1 | 79.9 | 114 | 18.2 | 12.3 | 25.8 | 35 |
| Urinary Bladder | 50.5 | 38.7 | 64.4 | 77 | 7.2 | 3.7 | 12.6 | 13 |
| Kidney and Renal Pelvis | 14.2 | 9.5 | 20.4 | 36 | 10.9 | 6.6 | 17.0 | 22 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pleura | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Bones and Joints | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Breast | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Ovary | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Vagina | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Vulva | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Male Genital System | 142.5 | 58.9 | 281.7 | 10 | 0.0 | 0.0 | 48.9 | 0 |
| Prostate | 142.5 | 58.9 | 281.7 | 10 | 0.0 | 0.0 | 48.9 | 0 |
| Testis | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Penis | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Ureter | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Alaska Native/American Indian | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Pancreas | 0.0 | 0.0 | 10.6 | 0 | 20.9 | 8.3 | 42.1 | 7 |
| Retroperitoneum | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 93.3 | 61.1 | 135.2 | 28 | 78.9 | 52.8 | 112.7 | 30 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 87.1 | 56.1 | 127.9 | 26 | 73.7 | 48.5 | 106.6 | 28 |
| Pleura | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Breast | 0.0 | 0.0 | 10.6 | 0 | 105.6 | 77.0 | 141.1 | 47 |
| Female Genital System | 0.0 | 0.0 | 10.6 | 0 | 63.1 | 41.7 | 91.4 | 29 |
| Cervix Uteri | 0.0 | 0.0 | 10.6 | 0 | 16.1 | 7.3 | 31.4 | 9 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 10.6 | 0 | 29.1 | 14.3 | 51.8 | 11 |
| Corpus Uteri | 0.0 | 0.0 | 10.6 | 0 | 29.1 | 14.3 | 51.8 | 11 |
| Uterus, NOS | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Ovary | 0.0 | 0.0 | 10.6 | 0 | 13.7 | 5.5 | 28.6 | 7 |
| Vagina | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Vulva | 0.0 | 0.0 | 10.6 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Male Genital System | 55.9 | 34.1 | 86.1 | 22 | 0.0 | 0.0 | 8.2 | 0 |
| Prostate | 41.6 | 23.3 | 68.4 | 16 | 0.0 | 0.0 | 8.2 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Urinary System | 33.9 | 16.9 | 60.0 | 12 | 14.1 | 5.0 | 30.8 | 6 |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 24.7 | 10.2 | 48.9 | 8 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Respiratory System | 69.7 | 25.6 | 145.2 | 7 | 53.9 | 22.8 | 105.3 | 8 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Lung and Bronchus | 69.7 | 25.6 | 145.2 | 7 | 46.5 | 18.3 | 95.2 | 7 |
| Pleura | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Bones and Joints | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Breast | 0.0 | 0.0 | 30.2 | 0 | 102.4 | 60.1 | 162.9 | 19 |
| Female Genital System | 0.0 | 0.0 | 30.2 | 0 | 52.6 | 24.2 | 99.4 | 10 |
| Cervix Uteri | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Ovary | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Male Genital System | 140.2 | 72.9 | 238.0 | 14 | 0.0 | 0.0 | 21.7 | 0 |
| Prostate | 125.2 | 61.2 | 220.5 | 12 | 0.0 | 0.0 | 21.7 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Penis | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Brain and Other Nervous System | 5.1 | 2.2 | 9.9 | 10 | 4.1 | 1.8 | 8.2 | 9 |
| Brain | 4.8 | 2.0 | 9.6 | 9 | 3.8 | 1.5 | 7.8 | 8 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 7.0 | 3.7 | 11.9 | 16 | 16.4 | 11.7 | 22.5 | 42 |
| Thyroid | 6.2 | 3.1 | 11.0 | 14 | 16.1 | 11.4 | 22.1 | 41 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 27.3 | 19.9 | 36.5 | 57 | 18.4 | 12.8 | 25.5 | 40 |
| Hodgkin Lymphoma | 5.2 | 2.6 | 9.3 | 13 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 5.2 | 2.6 | 9.3 | 13 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Non-Hodgkin Lymphoma | 22.2 | 15.4 | 30.8 | 44 | 17.5 | 12.0 | 24.6 | 38 |
| NHL - Nodal | 17.3 | 11.3 | 25.0 | 34 | 11.7 | 7.5 | 17.4 | 28 |
| NHL - Extranodal | 4.9 | 2.1 | 9.6 | 10 | 5.8 | 2.7 | 10.8 | 10 |
| Myeloma | 3.6 | 1.5 | 7.3 | 9 | 4.0 | 1.4 | 8.6 | 6 |
| Leukemia | 17.9 | 11.6 | 26.2 | 31 | 12.7 | 7.7 | 19.5 | 22 |
| Lymphocytic Leukemia | 10.0 | 5.8 | 16.0 | 20 | 6.6 | 3.2 | 11.9 | 11 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 7.1 | 3.6 | 12.5 | 13 | 5.4 | 2.2 | 10.5 | 8 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Myeloid and Monocytic Leukemia | 6.5 | 2.9 | 12.3 | 10 | 5.8 | 2.6 | 10.8 | 10 |
| Acute Myeloid Leukemia | 5.3 | 2.0 | 10.9 | 7 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 2.1 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 1.9 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | 0.0 | 0.0 | 2.1 | 0 | 0.0 | 0.0 | 1.9 | 0 |
| Miscellaneous | 16.2 | 10.0 | 24.6 | 27 | 10.0 | 5.6 | 16.2 | 17 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Lymphoma | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 64.6 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Mesothelioma | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 48.9 | 0 |
| Miscellaneous | 0.0 | 0.0 | 64.6 | 0 | 0.0 | 0.0 | 48.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Alaska Native/American Indian | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Lymphoma | 16.6 | 5.6 | 37.1 | 6 | 26.0 | 12.6 | 46.9 | 11 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 26.0 | 12.6 | 46.9 | 11 |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 17.6 | 6.8 | 36.3 | 7 |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 15.3 | 5.6 | 33.6 | 7 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 8.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 10.6 | 0 | 0.0 | 0.0 | 8.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2006, by Race and Sex

2006

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Urinary Organs | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Eye and Orbit | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Endocrine System | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 30.2 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.7 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Mesothelioma | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 30.2 | 0 | 0.0 | 0.0 | 21.7 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.